

## NR2B–NMDA receptor mediated modulation of the tyrosine phosphatase STEP regulates glutamate induced neuronal cell death

Ranjana Poddar,<sup>1</sup> Ishani Deb,<sup>1</sup> Saibal Mukherjee and Surojit Paul

University of New Mexico Health Sciences Center, Department of Neurology, University of New Mexico, Albuquerque, New Mexico, USA

### Abstract

The present study examines the role of a neuron-specific tyrosine phosphatase (STEP, striatal-enriched tyrosine phosphatase) in excitotoxic cell death. Our findings demonstrate that p38 MAPK, a stress-activated kinase that is known to play a role in the etiology of excitotoxic cell death is a substrate of STEP. Glutamate-mediated NMDA receptor stimulation leads to rapid but transient activation of p38 MAPK, which is primarily dependent on NR2A–NMDA receptor activation. Conversely, activation of NR2B–NMDA receptors leads to dephosphorylation and subsequent activation of STEP, which in turn leads to inactivation of p38 MAPK. Thus, during transient NMDA receptor stimulation, increases in STEP activity appears to limit the duration of activation of p38 MAPK and improves neuronal survival. However, if NR2B–NMDA recep-

tor stimulation is sustained, protective effects of STEP activation are lost, as these stimuli cause significant degradation of active STEP, leading to secondary activation of p38 MAPK. Consistent with this observation, a cell transducible TAT–STEP peptide that constitutively binds to p38 MAPK attenuated neuronal cell death caused by sustained NMDA receptor stimulation. The findings imply that the activation and levels of STEP are dependent on the duration and magnitude of NR2B–NMDA receptor stimulation and STEP serves as a modulator of NMDA receptor dependent neuronal injury, through its regulation of p38 MAPK.

**Keywords:** glutamate, NR2A, NR2B, p38 mitogen activated protein kinase, STEP, tyrosine phosphatase.

*J. Neurochem.* (2010) **115**, 1350–1362.

Excitotoxic glutamate receptor activation is implicated in neuro-degeneration following ischemic insults, post-traumatic lesions and a range of other neurological disorders (Lipton and Rosenberg 1994; Arundine and Tymianski 2004). In pathological conditions excessive release of the neurotransmitter glutamate leads to sustained activation of NMDA-type glutamate receptors and a deleterious cascade of intracellular events that include  $Ca^{2+}$  overload and activation of effectors such as p38 MAPK. Recent findings suggest the existence of two functionally distinct pools of NMDA receptors defined by the presence of NR2A (NR2A–NMDAR) or NR2B (NR2B–NMDAR) containing subunits (Luo *et al.* 1997; Tovar and Westbrook 1999; Kohr 2006). NR2B–NMDARs are generally thought to be involved in triggering cell death and subsequent brain damage following an excitotoxic insult (Liu *et al.* 2007; Tu *et al.* 2010). However, pharmaceutical application of NR2B–NMDAR antagonists have been unsuccessful in clinical trials, as they interfere with physiological functions of these receptors, in addition to blocking excitotoxic cascades (Rodrigues *et al.*

2001; Zhao *et al.* 2005; Walker and Davis 2008). An alternative approach that may improve treatment of excitotoxic insults involves targeting of components of intracellular signaling cascades downstream of NMDA receptor signaling.

The p38 MAPK pathway has been shown to be involved in mediating the pathogenesis of glutamate excitotoxicity (Barone *et al.* 2001a,b). While the role of NMDA receptors in the activation of p38 MAPK is well established (Mukherjee *et al.* 1999; Waxman and Lynch 2005), the precise

Received May 20, 2010; revised manuscript received August 20, 2010; accepted September 27, 2010.

Address correspondence and reprint requests to Surojit Paul, University of New Mexico Health Sciences Center, Department of Neurology, 1, University of New Mexico, Albuquerque, NM 87131, USA. E-mail: spaul@salud.unm.edu

<sup>1</sup>These authors contributed equally to this study.

**Abbreviations used:** APV, DL-2-amino-5-phosphonopentanoic acid; ERK, extracellular regulated kinase; GST, Glutathione S-transferase; KIM, kinase interacting motif; PBS, phosphate-buffered saline; PKA, protein kinase A; STEP, striatal-enriched phosphatase; WT, wild-type.

intracellular mechanisms that regulate the duration and magnitude of p38 MAPK are not clearly understood. A goal of the present work was to determine whether the p38 MAPK activation is significantly regulated by the neuron-specific tyrosine phosphatase, STEP (striatal-enriched tyrosine phosphatase) and whether this could be targeted to limit excitotoxic injury.

Striatal-enriched tyrosine phosphatase, also known as PTPN5, is expressed specifically in neurons of the striatum, neo-cortex and hippocampus (Boulanger *et al.* 1995). STEP<sub>61</sub> and STEP<sub>46</sub>, the two STEP isoforms (Bult *et al.* 1997) contain a highly conserved substrate-binding domain termed as the kinase interacting motif (KIM) domain (Pulido *et al.* 1998). Phosphorylation of a critical serine residue within the KIM domain is mediated through dopamine/D<sub>1</sub> receptor dependent activation of the protein kinase A (PKA) pathway (Paul *et al.* 2000). Dephosphorylation of this residue by Ca<sup>2+</sup> dependent phosphatase calcineurin, following glutamate/NMDA receptor stimulation, renders STEP active in terms of its ability to bind to its substrates (Paul *et al.* 2003). Active STEP can bind to and inhibit the activity of ERK (extracellular regulated kinase 1/2) MAPK (Paul *et al.* 2007).

The aim of the present study was to investigate the functional significance of STEP's activation following excitotoxic NMDAR stimulation. Our findings show that NR2B-NMDAR stimulation leads to activation of STEP. Active STEP can contribute to neuronal cell survival by down-regulating p38 MAPK activity. However, sustained NR2B-NMDAR stimulation results in the degradation of active STEP, and relieves its inhibitory effect on p38 MAPK and results in cell death.

## Materials and methods

### Materials and reagents

Pregnant female and adult (250 g) Sprague–Dawley rats were from Harlan Laboratories (Livermore, CA, USA). Antibodies used were as follows: polyclonal anti-p38, rabbit monoclonal anti-phospho-p38 (T<sup>P</sup>EY<sup>P</sup> and TEY<sup>P</sup>) from Cell Signaling (Danvers, MA, USA), monoclonal anti-myc from Santa Cruz Biotechnology (Santa Cruz, CA, USA), monoclonal anti-Flag and polyclonal anti-tubulin from Sigma-Aldrich (St. Louis, MO, USA), polyclonal anti-MAP2 from Chemicon, Temecula, CA, USA, monoclonal anti-V5 and anti-V5-HRP from Invitrogen (Carlsbad, CA, USA), monoclonal anti-STEP from Novus Biologicals (Littleton, CO, USA). All secondary antibodies were from Cell Signaling. All tissue culture reagents were obtained from Invitrogen. All other reagents were from Sigma-Aldrich. Approval for animal experiments was given by the University of New Mexico, Health Sciences Center, Institutional Animal Care and Use Committee.

### Plasmids and transfection

STEP<sub>46</sub> cDNA was constructed in mammalian expression vector pcDNA 3.1 encoding a C-terminal V5 and His tag (Invitrogen). Point mutants were generated by PCR-based site-directed

mutagenesis using Pfu Turbo DNA polymerase (Stratagene, La Jolla, CA, USA) according to the manufacturer's protocol. All mutations were verified by nucleotide sequencing. The p38 $\alpha$ ,  $\beta$  and  $\gamma$  cDNAs with a FLAG tag at the C-terminal were gifts from Dr. Anton Bennett, Yale University.

### *In vitro* phosphorylation of STEP and tyrosine phosphatase assay

Glutathione S-transferase (GST), GST-STEP<sub>46</sub> and GST-STEP<sub>46</sub> C300S fusion proteins were generated as described previously (Paul *et al.* 2000). For *in vitro* binding studies GST fusion proteins were incubated with p38 $\alpha$  MAPK recombinant protein (EMD Biosciences, San Diego, CA, USA) overnight at 4°C in a buffer containing 50 mM Tris-HCl, pH 7.0, 100 mM NaCl, 5 mM dithiothreitol, 0.01% Brij 35 and 2 mM EDTA. Samples were then incubated with Glutathione-sepharose beads (2 h) and washed extensively. Bound proteins were eluted using sodium dodecyl sulfate-sample buffer and processed for immunoblot analysis. *In vitro* phosphorylation of GST-STEP<sub>46</sub> fusion protein by PKA was done as described earlier (Paul *et al.* 2003). Phosphorylated and non-phosphorylated STEP was incubated with active p38 $\alpha$  MAPK recombinant protein for specified time periods (0, 5, 7.5 and 10 min). The reactions were stopped by addition of sodium dodecyl sulfate-sample buffer and the samples were processed for immunoblot analysis.

### Affinity column chromatography

STEP<sub>46</sub> C300S cDNA, a substrate-trapping variant of STEP<sub>46</sub>, was cloned into the *E. Coli* expression vector pET104.1, with a six-histidine tag at the C-terminal (Invitrogen). The vector provides a biotin conjugation site at the N-terminus of the protein to allow *in vivo* biotinylation of the over-expressed protein by endogenous *E. Coli* biotin ligases. The purified biotinylated protein was coupled to streptavidin agarose beads (Thermo Scientific, Rockford, IL, USA). Using striatal homogenates from rat brain substrate-trapping experiments were performed as described before (Nguyen *et al.* 2002). Bound proteins were eluted with 0.1 M glycine, pH 2.8 and processed for immunoblot analysis.

### Purification of TAT-STEP-myc fusion protein

STEP<sub>46</sub>  $\Delta$ PTP cDNA (STEP<sub>46</sub> 1–107) that lacks the C-terminal phosphatase domain was subcloned in a pTrc-His-myc-TOPO expression vector (Invitrogen). To render the peptide cell permeable an 11 amino acid peptide called TAT was inserted at the N-terminal of the STEP<sub>46</sub>  $\Delta$ PTP cDNA (Paul *et al.* 2007). A point mutation was introduced at the serine residue within the KIM domain (S49A) by site-directed mutagenesis. STEP<sub>46</sub>  $\Delta$ PTP was expressed in *E. Coli* and the fusion protein was purified using BD-Talon resin (BD Biosciences, Bedford, MA, USA).

### Cell culture and stimulation

Primary neuronal cultures were obtained from 16- to 17-day-old rat embryos as described previously (Paul *et al.* 2003). Briefly, the striatum and the adjoining cortex was dissected, the tissue dissociated mechanically and resuspended in Dulbecco's modified Eagle's medium/F-12 (1 : 1)-containing 5% fetal calf serum. Cells ( $6 \times 10^6$  cells/dish) were plated on poly-D-lysine-coated tissue culture dishes (BD Biosciences) and grown for 12–14 days at 37°C in a humidified atmosphere (95 : 5% air : CO<sub>2</sub> mixture). For

receptor stimulation cells were treated with glutamate or NMDA for the indicated times and then processed for immunoprecipitation or immunoblot analysis as described earlier (Paul *et al.* 2003). Glycine concentration in the medium during glutamate treatment was 1  $\mu$ M. In some experiments, DL-2-amino-5-phosphonopentanoic acid (APV) or 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX) were added 5 min after the addition of glutamate; ifenprodil and Ro 25-6981 were added 10 min before addition of glutamate. Some cultures were treated with the TAT-STEP-myc peptide (4  $\mu$ M) or SB-203580 (2.6  $\mu$ M) prior to or along with glutamate treatment. Densitometric analysis of immunoblots was performed using Image J software (NIH, Bethesda, MD, USA). Statistical comparison was carried out using one-way analysis of variance (ANOVA, Bonferroni's multiple comparison test) and differences were considered significant when  $p < 0.05$ .

#### Immunocytochemistry and Hoechst DNA staining

Hela cells transfected with STEP<sub>46</sub> cDNA or neuron cultures grown for 12–14 days were processed for immunocytochemistry as described earlier (Paul *et al.* 2003). Hela cells were immunostained with anti-phospho-p38 (T<sup>P</sup>EY<sup>P</sup>, 1 : 100) and/or anti-STEP (1 : 250) antibodies. Neuronal cells were immunostained with anti-myc (1 : 100), anti-MAP2 (1 : 250) or anti-phospho p38 (T<sup>P</sup>EY<sup>P</sup>, 1 : 100) antibodies. Alexa Fluor 488 conjugated anti-mouse IgG or Cy3 conjugated anti-rabbit IgG were used as secondary antibodies (1 : 250 each). In some cases, neurons were subjected to staining with Hoechst 33342 dye for 15 min to assess nuclear damage. Imaging was performed with a Zeiss Axiovert 200M fluorescence microscope with attached AxioCam CCD camera using 20 $\times$  objective lens (Carl Zeiss, Thornwood, NY, USA). To quantitatively assess the percentage of pyknotic nuclei a total of 1000 cells were counted for each set of experiments. Mean  $\pm$  SEM ( $n = 3$ ) were used for statistical comparison using ANOVA (Bonferroni's multiple comparison test). Differences were considered significant when  $p < 0.05$ .

#### Stereotactic surgery and immunohistochemistry

For intra-striatal injection of glutamate adult male rats were anesthetized with 2% isoflurane in 70% nitrous oxide and 30% oxygen. A burr hole was drilled and a micropipette was lowered into the right striatum using the following coordinates (Paxinos and Watson 1986): 5.0 mm lateral to bregma and 5.5 mm ventral to the skull surface. 1  $\mu$ L of TAT-STEP-myc peptide or phosphate-buffered saline (PBS) was infused into the right striatum over a time period of 2 min. The injector was left in place for an additional 2 min to minimize back-flux of the solution. After injector withdrawal, the wound was sutured and the rats returned to their cages. In some animals, 30 nmol of glutamate in 1  $\mu$ L PBS or PBS alone (control) was infused into the striatum 30 min after infusion of TAT-STEP-myc peptide or PBS. At the specified time points after infusion rats were anesthetized, perfused intracardially with 4% paraformaldehyde and processed for cryosectioning followed by immunohistochemistry with anti-myc antibody (1 : 100) (Paul *et al.* 2007). Sections were then treated with Hoechst 33342 dye for DNA stain, mounted using vectashield and analyzed using fluorescent microscopy. For assessment of nuclear damage, the percentage of pyknotic nuclei was estimated in five different fields within the striatum of each section immediately below the site of needle injection. For sections obtained from TAT-STEP-myc peptide

injected animals the percentage of cells positively stained with anti-myc antibody within the same fields were also determined. Each experiment was repeated at least three times. Statistical comparison was carried out using ANOVA (Bonferroni's multiple comparison test) and differences were considered significant when  $p < 0.05$ .

## Results

### Glutamate–NMDA receptor mediated regulation of p38 MAPK in neurons


We first determined the temporal profile of p38 MAPK phosphorylation following an excitotoxic insult. Neuronal cultures (12–14 days *in vitro*) were treated with 100  $\mu$ M glutamate for varying times and cell lysates were analyzed by immunoblotting with a phospho-specific antibody that recognizes p38 only when phosphorylated at both the regulatory threonine and tyrosine residues (T<sup>P</sup>EY<sup>P</sup>-p38). Our results show a transient increase in phosphorylation of p38 MAPK within 2–5 min of glutamate stimulation. This was followed by a time-dependent decrease in p38 phosphorylation, which returned to near basal levels by 30 min (Fig. 1a). Total p38 MAPK level remained unaltered by this treatment. Treatment with 100  $\mu$ M NMDA caused a similar transient activation of p38 MAPK (Fig. 1b) as was seen with glutamate suggesting that NMDAR activation was required for phosphorylation of p38 MAPK.

To explore the contribution of NMDARs in the dephosphorylation of p38 MAPK, a selective NMDA receptor antagonist (APV, 200  $\mu$ M), was applied 5 min after incubation with glutamate. The delayed treatment with APV inhibited the glutamate-induced dephosphorylation of p38 MAPK (Fig. 1c). However, a similar treatment with 6-cyano-7-nitroquinoxaline-2,3-dione (20  $\mu$ M), a non-NMDA receptor antagonist failed to prevent the glutamate-induced dephosphorylation of p38 MAPK (Fig. 1d). These observations in conjunction with previous studies (Waxman and Lynch 2005) suggest that both phosphorylation and dephosphorylation of p38 MAPK in neurons require NMDA receptor activation.

We next examined the roles of NR2A- and NR2B-NMDARs in the dephosphorylation of p38 MAPK. Neurons were treated with 100  $\mu$ M glutamate in the presence or absence of selective NR2B-NMDAR antagonists, ifenprodil and Ro 25-6981. As shown in Fig. 1(e and f) these antagonists had no effect on the initial phosphorylation of p38 MAPK by 5 min, however, the subsequent dephosphorylation of p38 MAPK (measured at 30 min) was significantly reduced, suggesting that NR2B-NMDAR stimulation are required for the dephosphorylation of p38 MAPK.

### NR2B–NMDAR mediated activation of STEP followed by its proteolytic cleavage

A potential candidate that might mediate the NR2B-NMDAR dependent dephosphorylation of p38 MAPK is the tyrosine




**Fig. 1** Role of NR2B-NMDARs in glutamate-NMDAR induced transient phosphorylation of p38 MAPK. Neuron cultures were treated with (a) glutamate (100 μM) or (b) NMDA (100 μM) for the specified times. (c–d) Neurons were treated with glutamate for the specified times. For lanes 4 and 5, after treatment with glutamate for 5 min, APV (c) or CNQX (d) was added and the incubation was continued for the specified times. (e, f) Neurons were treated with 100 μM glutamate for specified times in the presence or absence of ifenprodil (e) or Ro 25-6981 (f). Equal amount of protein from each sample was processed for immunoblot analysis using anti-phospho p38 (T<sup>P</sup>EY<sup>P</sup>-p38, upper panel) and p38 (lower panel) antibodies. Quantification of phosphorylated p38 MAPK by computer-assisted densitometry and Image J analysis is shown below each immunoblot. Values are mean ± SEM (*n* ≥ 3). \*Significant difference from 5-min glutamate or NMDA treatment (*p* < 0.001). #Significant difference from 30-min glutamate treatment (*p* < 0.001).

phosphatase STEP. Under basal condition STEP is phosphorylated at the PKA site in the KIM domain (Ser221 in STEP<sub>61</sub> and Ser49 in STEP<sub>46</sub>) (Paul *et al.* 2000). Stimulation of NMDARs by glutamate leads to time-dependent dephosphorylation and activation of STEP, which begins at 5 min and leads to complete dephosphorylation by 30 min. Using a phospho-specific antibody (against the PKA site) it has also been shown that dephosphorylation of STEP results in downward shift in mobility of the STEP band (Paul *et al.* 2003). Consistent with these earlier findings incubation of neurons with glutamate resulted in a marked decrease in the upper band and the appearance of a predominant lower band, indicating that STEP<sub>61</sub> is rapidly dephosphorylated following glutamate treatment (Fig. 2a and b, lane 2). Pre-incubation with either Ro 25-6981 or ifenprodil blocked the downward

shift in the mobility of the upper band, supporting a role for NR2B-NMDAR in the dephosphorylation of STEP<sub>61</sub> (Fig. 2a and b, lane 3).

We also examined the temporal profile of STEP's activation following treatment with glutamate (100 μM, 30 min), by immunoblot analysis with anti-STEP antibody. Figure 2(c) shows that 30-min after termination of glutamate treatment STEP was subjected to proteolytic cleavage resulting in the appearance of a cleaved product of ~33 kDa (STEP<sub>33</sub>), which increased over time (~5-fold 1 h). Analyzing the same samples with anti-T<sup>P</sup>EY<sup>P</sup> antibody showed an increase in phosphorylation of p38 MAPK 0.5–1 h after termination of glutamate exposure (Fig. 2d). To determine the involvement of NR2B-NMDAR in the cleavage of STEP neurons were incubated with Ro 25–



**Fig. 2** Glutamate/NMDAR induced dephosphorylation and subsequent cleavage of STEP involves NR2B-NMDARs. Neuronal cultures were treated with (a–b) 100  $\mu$ M glutamate for 30 min in the presence or absence of Ro 25-6981 (a) or ifenprodil (b); (c–d) 100  $\mu$ M glutamate for 30 min and maintained in its original medium for the specified times (post-glutamate time); (e) 100  $\mu$ M glutamate for 30 min in the presence or absence of Ro 25-6981 and maintained in its original medium for the specified time (post-glutamate time). Extracts with equal amount of protein were analyzed from each sample with anti-STEP antibody (a, b, c; upper and middle panel of e). In some samples, phosphorylation of p38 was analyzed using anti-phospho p38 antibody (T<sup>P</sup>EY<sup>P</sup>-p38; upper panel, d). Total p38 or tubulin was analyzed by re-probing the membrane with either anti-p38 (lower panel, d) or anti-tubulin antibody (lower panel, e). Quantification of (a–b) phosphorylated (upper band) and non-phosphorylated (lower band) STEP was done using computer assisted densitometry and Image J analysis and data presented as percentage of the total; (c, e) extent of proteolytic cleavage of STEP and (d) phosphorylation of p38 MAPK was done using computer assisted densitometry and Image J analysis. Values are mean  $\pm$  SEM ( $n \geq 3$ ). \*Significant difference from control ( $p < 0.001$ ). #Significant difference from 30-min or 1-h glutamate treatment ( $p < 0.001$ ).

6981 prior to treatment with glutamate (100  $\mu$ M glutamate, 30 min). Pre-incubation with Ro 25-6981 blocked the dephosphorylation as well as the proteolytic cleavage of STEP<sub>61</sub> (Fig. 2e). These results imply that glutamate-mediated NR2B-NMDAR stimulation leads to initial activation of STEP, and then a progressive decrease in STEP levels over time in the post-glutamate period. When taken together with results above (Fig. 1), these observations raise the possibility that active STEP may play an initial role in dephosphorylation of p38 MAPK and may serve as a key regulator of p38 MAPK activation in excitotoxicity. However, degradation of STEP may lead to the secondary activation of p38 MAPK.


### p38 MAPK is a substrate of STEP

To determine whether p38 MAPK is a potential substrate of STEP, striatal homogenates from rat brain were subjected to affinity purification with a substrate-trapping mutant of STEP (STEP<sub>46</sub> C300S). Immunoblot analysis of the eluant showed that p38 MAPK was part of the protein complex bound to STEP (Fig. 3a). To determine if STEP can directly interact with p38 MAPK, purified GST-STEP<sub>46</sub> or GST-STEP<sub>46</sub> C300S protein coupled to glutathione sepharose beads was incubated with a p38 MAPK fusion protein. Immunoblot analysis revealed that both variants of STEP were able to pull down p38 MAPK (Fig. 3b). To test the ability of active STEP to dephosphorylate p38 MAPK *in vitro*, purified GST-STEP<sub>46</sub>

was incubated with phosphorylated p38 MAPK fusion protein (active form) for varying time periods. Immunoblot analysis with an antibody that recognizes p38 only when phosphorylated at the tyrosine residue (TEY<sup>P</sup> p38) revealed a time-dependent decrease in tyrosine phosphorylation of p38 MAPK (Fig. 3c, upper panel). The ability of phosphorylated STEP (phosphorylated at the PKA site in the KIM domain) to dephosphorylate active p38 MAPK was also measured. The

data shows that phosphorylated STEP fails to dephosphorylate p38 MAPK (Fig. 3c, lower panel). These results suggest that p38 MAPK is a substrate of STEP and phosphorylation of STEP at the serine residue in the KIM domain inhibits the interaction between STEP and p38 MAPK.

To further clarify the mechanism by which STEP regulates p38 MAPK in intact cells V5-STEP<sub>46</sub> wild-type (WT) or one of its mutants that cannot be phosphorylated endogenously at



**Fig. 3** p38 MAPK is a substrate of STEP. (a) Rat brain homogenates were passed through STEP<sub>46</sub> C300S affinity column. Eluants were processed for immunoblot analysis using anti-p38 MAPK antibody (SM, starting material; E, eluant). (b) GST, GST-STEP<sub>46</sub> or GST-STEP<sub>46</sub> C300S fusion proteins were incubated with recombinant p38 MAPK protein. Bound proteins were extracted and processed for immunoblot analysis using anti-p38 MAPK (upper panel) antibody. Blots were re-probed with anti-GST (lower panel) antibody. (c) GST-STEP<sub>46</sub> or GST-STEP<sub>46</sub> phosphorylated with PKA were incubated with active p38 $\alpha$  MAPK (phosphorylated at the regulatory threonine and tyrosine residues) for the specified times. Samples were analyzed using a phospho-specific antibody that recognizes p38 MAPK when it is phosphorylated at the regulatory tyrosine residue

(TEY<sup>P</sup>-p38). Blots were re-probed with anti-p38 MAPK antibody. (d–e) HeLa cells were transfected with (d) V5-STEP<sub>46</sub> or one of its mutants (S49A and S49A/C300S); or (e) V5-STEP<sub>46</sub> S49A and Flag-tagged p38 $\alpha$ ,  $\beta$ ,  $\gamma$  or  $\delta$ . STEP was immunoprecipitated with anti-V5 antibody. Samples were analyzed either with anti-p38 (upper panel, d) or anti-Flag antibody (upper panel, e). Blots were re-probed with anti-V5-HRP antibody (lower panel, d–e). In (e) total lysate was analyzed using anti-Flag antibody to demonstrate expression of all p38 isoforms in HeLa cells. (f–g) Untransfected HeLa cells (f) and HeLa cell transfected with V5-STEP<sub>46</sub> or V5-STEP<sub>46</sub>-S49A (g) were treated with 0.3 M sorbitol for 15 min and processed for immunocytochemical staining with anti-phospho p38 (T<sup>P</sup>EY<sup>P</sup>-p38; f, g) and anti-V5-HRP (g) antibodies.

the KIM domain site (S49A or S49A/C300S) were transfected in HeLa cells. Immunoprecipitation of STEP using anti-V5 antibody followed by immunoblot analysis using anti-p38 antibody showed that both V5-STEP<sub>46</sub> S49A (dephosphorylated form of STEP) and V5-STEP<sub>46</sub> S49A/C300S (dephosphorylated and catalytically inactive mutant) were able to pull down endogenous p38 MAPK (Fig. 3d). In contrast the phosphorylated form (V5-STEP<sub>46</sub> WT) failed to bind to p38 MAPK. This suggests that in intact cells only the dephosphorylated form of STEP can interact with p38 MAPK.

The family of p38 MAPKs includes p38 $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  isoforms, although only p38 $\alpha$  and  $\beta$ , are expressed in the brain (Kumar *et al.* 2003). To determine whether STEP can interact with all four isoforms, HeLa cells were co-transfected with V5-STEP<sub>46</sub> S49A and either p38 $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  cDNA. Only p38 $\alpha$  and  $\beta$  immunoprecipitated with STEP (Fig. 3e) suggesting a role of STEP in regulating the physiological function of these two isoforms in neurons.


Activation and subsequent nuclear translocation of p38 MAPK has been implicated in glutamate-NMDA receptor induced neuronal cell death (Waxman and Lynch 2005). To examine the role of STEP in regulating the nuclear translocation of active p38 MAPK, HeLa cells were transfected with V5-STEP<sub>46</sub> WT or the S49A mutant and then treated with sorbitol. Consistent with earlier studies (Ingram *et al.* 2000) hyper-osmotic stress generated with sorbitol resulted in phosphorylation and subsequent nuclear translocation of active p38 MAPK (Fig. 3f). In the presence of V5-STEP<sub>46</sub> WT, sorbitol-mediated nuclear translocation of p38 MAPK was not affected. In contrast, expression of V5-STEP<sub>46</sub> S49A abolished nuclear translocation of active p38 MAPK (Fig. 3g). These results establish that only the dephosphorylated and active form of STEP can block phosphorylation and nuclear translocation of p38 MAPK.

The above findings show that p38 MAPK is inhibited by STEP and STEP's activation following glutamate/NMDA receptor stimulation correlates strongly with the inactivation of p38 MAPK. While these results suggest a neuroprotective role for active STEP, it is clear that under most excitotoxicity paradigms STEP activation is not sufficient to prevent neuronal injury. This is illustrated in Figure S1, where delayed neuronal death is observed following exposure of neuron cultures to 100  $\mu$ M glutamate for 30 min, as has been described in many prior reports (Kawasaki *et al.* 1997; Barone *et al.* 2001a). This delayed neuronal cell death was blocked by pre-incubation with SB-203580 a selective inhibitor of p38 MAPK. This confirms a central role of p38 MAPK in the excitotoxic cascade and emphasizes that any inhibitory effects of STEP were not sufficient to block p38 MAPK activation (Figure S1). This is possibly because of the fact that sustained stimulation of NMDARs led to proteolytic cleavage of active STEP resulting in secondary activation of p38 MAPK.

### TAT-STEP attenuates glutamate/NMDA receptor induced neuronal cell death


To directly test the hypothesis that interaction of STEP with p38 MAPK regulates neuronal injury we generated a cell permeable TAT-STEP-myc peptide (TAT-STEP<sub>46</sub> S49A  $\Delta$ PTP-myc, see 'Materials & Methods') that can constitutively bind to p38 MAPK. We first determined the temporal profile of intracellular uptake of the TAT-STEP-myc peptide. Neuronal cultures were incubated with the TAT-STEP-myc peptide for varying time periods (0, 5, 15, 30 or 120 min). Immunocytochemical analysis with anti-myc antibody showed that the peptide was detectable 15 min after application that peaked at 30 min and was present for the rest of the time studied (Fig. 4a). We then determined whether the TAT-STEP-myc peptide could bind to endogenous p38 MAPK. Neuronal cultures were incubated with the peptide for 30 min followed by immunoprecipitation with anti-myc antibody. Immunoblotting with anti-p38 antibody showed that the TAT-STEP-myc peptide forms a stable complex with p38 MAPK (Fig. 4b). We next examined whether pre-incubation with TAT-STEP-myc peptide or its co-application along with glutamate can block the initial phosphorylation and nuclear translocation of p38 MAPK at 5 min. Stimulation with glutamate (100  $\mu$ M, 5 min) alone led to phosphorylation (Fig. 4c and d, upper panel, lane 2) and nuclear translocation of p38 MAPK (Fig. 4e, upper panel). Pre-incubation of neurons with TAT-STEP-myc peptide (30 min) prior to glutamate treatment blocked the phosphorylation (Fig. 4c, upper panel, lane 3) and nuclear translocation of p38 MAPK at 5 min (Fig. 4e, middle row). When the peptide was co-administered along with glutamate, it failed to block the initial phosphorylation (Fig. 4d, upper panel, lane 3) and nuclear translocation (Fig. 4e, lower panel) of p38 MAPK by 5 min. Finally, we examined if co-application of TAT-STEP-myc peptide along with glutamate can block the delayed phosphorylation and nuclear translocation of p38 MAPK at the post-glutamate period, between 0.5 and 1 h. Neurons were incubated with glutamate (100  $\mu$ M) for 30 min in the presence or absence of TAT-STEP-myc peptide and then processed for immunoblotting and immunocytochemical analysis 30 min after termination of glutamate treatment. Stimulation with glutamate alone led to phosphorylation (Fig. 4f, upper panel, lane 2) and nuclear translocation of p38 MAPK (Fig. 4g, upper row). Co-application of TAT-STEP-myc peptide blocked the phosphorylation (Fig. 4f, upper panel, lane 3) and nuclear translocation (Fig. 4g, lower row) of p38 MAPK.

We then evaluated the role of the initial p38 MAPK activation by 5 min as well as the delayed p38 MAPK activation between 0.5 and 1 h (post-glutamate period) in neuronal injury following exposure to an excitotoxic dose of glutamate (100  $\mu$ M, 30 min). For these experiments the TAT-STEP-myc peptide was applied to neuronal cultures either 30 min prior to glutamate treatment (100  $\mu$ M, 30 min) or co-



**Fig. 4** TAT-STEP-myc peptide can block phosphorylation and nuclear translocation of p38 MAPK. (a) Neuronal cultures were incubated with the TAT-STEP-myc peptide for the specified times and processed for immunocytochemical staining with anti-myc antibody. (b) Neuronal cultures were incubated with TAT-STEP-myc peptide (30 min) followed by immunoprecipitation with anti-myc antibody. Samples were analyzed using anti phospho-p38 antibody ( $T^PEY^P$ -p38, upper panel). Blots were re-probed with anti-myc antibody (lower panel). (c–e) Neuronal cultures were treated with glutamate (100 μM, 5 min) in the absence or presence of TAT-STEP-myc peptide. The peptide was applied to neurons either 30 min prior to (c, e: middle panel) or along with glutamate (d, e: lower panel). (c–d) Cell lysates were analyzed

using anti-phospho-p38 antibody ( $T^PEY^P$ -p38, upper panel). Blots were re-probed with anti-p38 (c, d: middle panel) or anti-myc (c, d: lower panel) antibodies. (e) Cells were processed for immunocytochemical analysis with anti-phospho p38 antibody ( $T^PEY^P$ -p38) and DAPI. (f–g) Neuron cultures were treated with glutamate (100 μM, 30 min) in the absence or presence of TAT-STEP-myc peptide. The peptide was co-applied along with glutamate. 30 min after termination of glutamate treatment cells were processed either for (f) immunoblot analysis using anti-phospho-p38 ( $T^PEY^P$ -p38, upper panel), -p38 (middle panel) and -myc (lower panel) antibodies, respectively or (g) for immunocytochemical analysis with anti-phospho p38 antibody ( $T^PEY^P$ -p38) and DAPI.



**Fig. 5** TAT-STEP-myc peptide reduces glutamate-induced neuronal cell death. Neuronal cultures were treated with glutamate (100  $\mu$ M, 30 min) in the absence or presence of TAT-STEP-myc peptide or the corresponding TAT-myc control and then maintained in the original medium for another 20 h. TAT-STEP-myc peptide was applied to cells either 30 min prior to or along with glutamate. (a) Representative photomicrograph demonstrating the extent of cell death, by immunocytochemical analysis with anti-MAP2 antibody and Hoechst DNA staining. (b) Quantitative analysis of the percentage of neurons with pyknotic nuclei is represented as mean  $\pm$  SEM ( $n \geq 3$ ). \*Significant difference from 30-min glutamate treatment ( $p < 0.001$ ).

applied along with glutamate (100  $\mu$ M, 30 min). Twenty hours later, cells were processed for Hoechst DNA staining and MAP2 staining to test for cell viability. The representative photomicrographs and the corresponding mean data (Fig. 5a and b) demonstrate that  $16 \pm 5\%$  of the cells were apoptotic under basal conditions (no treatment). Following glutamate treatment  $79 \pm 7\%$  of the cells became apoptotic. Pre-incubation with TAT-myc peptide (30 min) alone failed to attenuate glutamate induced neuronal cell death. Pre-incubation with TAT-STEP-myc peptide (30 min) prior to glutamate treatment greatly reduced the number of apoptotic cells to  $26 \pm 4\%$ . Co-application of the TAT-STEP-myc peptide along with glutamate also reduced the number of apoptotic cells to  $30 \pm 4\%$ . This latter result is important when taken together with the results of the p38 MAPK activation above (Fig. 4). Thus, TAT-STEP-myc peptide when co-administered along with glutamate failed to block the initial p38 MAPK activation by 5 min but still abolished the delayed p38 MAPK activation and greatly reduced neuronal injury. This is consistent with our hypothesis that neuronal injury following exposure to an excitotoxic dose of glutamate is primarily because of delayed activation of p38 MAPK following the proteolytic cleavage of STEP.


#### TAT-STEP-myc attenuates cell death *in vivo*

We next investigated whether the TAT-STEP-myc peptide infused in the brain can prevent excitotoxic injury *in vivo*. This was tested in the well-established model of glutamate

microinjection into the striatum and evaluation of neuronal injury 24 h later (Iadecola *et al.* 2001; Haelewyn *et al.* 2008). In initial experiments, uptake of the TAT-STEP-myc peptide into striatal cells was confirmed following stereotactic injection into the right striatum (Fig. 6a and b). Coronal brain sections of the striatum taken at different time points (12 and 24 h) after infusion were examined immunohistochemically with anti-myc antibody. While no myc staining was observed in the contralateral striatum (left striatum), myc positive cells were detectable in the ipsilateral striatum (right striatum) infused with TAT-STEP-myc, at both the time points studied (Fig. 6c). In subsequent experiments, animals were stereotactically injected with glutamate (30 nmol) in the presence or absence of TAT-STEP-myc peptide. 24 h later coronal brain sections of the striatum were processed for immunohistochemical staining with anti-myc antibody and Hoechst DNA staining. The representative photomicrograph and the corresponding mean data (Fig. 6d) show a significant reduction in cell death in the presence of the peptide. Quantitative analysis further showed that  $93 \pm 1.1\%$  of the cells that were transduced with TAT-STEP-myc peptide was protected against glutamate-induced cell death.

#### Discussion

A key finding of the present study is that NR2B-NMDARs play a critical role in neuronal survival through activation of the tyrosine phosphatase STEP that in turn can limit the



**Fig. 6** TAT-STEP-myc peptide reduces glutamate-induced cell death in the striatum. (a) Schematic representation of the area of infusion of TAT-STEP-myc peptide. (b–c) TAT-STEP-myc peptide was stereotactically injected into the striatum. Coronal sections were processed for immunohistochemical analysis with anti-myc antibody at the specified time periods. (b) Low magnification (5 $\times$ ) and (c) high magnification (20 $\times$ ) images. (d) Glutamate was injected

stereotactically into the striatum in the presence or absence of TAT-STEP-myc peptide. Striatal brain sections were processed for immunohistochemical analysis with anti-myc antibody and Hoechst DNA stain (left panel). Quantitative analysis of the percentage of neurons with pyknotic nuclei is represented as mean  $\pm$  SEM ( $n \geq 3$ , right panel). \*Significant difference from glutamate treated samples ( $p < 0.001$ ).

duration of activation of p38 MAPK. However, active STEP is susceptible to degradation on prolonged exposure to an excitotoxic insult leading to secondary activation of p38 MAPK. The ability of TAT-STEP-myc peptide to inhibit the delayed activation of p38 MAPK and attenuate cell death further suggests that activation of pro-apoptotic pathways are possibly downstream of the secondary p38 MAPK activation. It is also conceivable that in addition to p38 MAPK there may be other potential substrates that STEP could act upon to confer neuroprotection.

This study in conjunction with previous findings (Waxman and Lynch 2005) indicates that stimulation of NR2A-NMDARs leads to a rapid but transient activation of p38 MAPK. Previous studies have also shown that glutamate-mediated NR2A-NMDAR stimulation leads to transient activation of ERK MAPK (Kim *et al.* 2005). Such transient activation of ERK and p38 MAPK pathways under physiological conditions is thought to be involved in the induction of long-term potentiation and long-term depression, respectively (Bolshakov *et al.* 2000; Zhu *et al.* 2002; Gisabella *et al.* 2003; Thomas and Haganir 2004; Wang *et al.* 2004), two opposing forms of synaptic plasticity that are thought to contribute to learning and memory. The transient nature of p38 and ERK activation following NR2A-NMDAR stimulation suggests sequential activation of a stimulatory and an

inhibitory pathway. The inhibitory pathway involves the tyrosine phosphatase STEP. Glutamate-mediated NMDA receptor stimulation leads to delayed activation of STEP via  $Ca^{2+}$ -dependent phosphatase calcineurin (Paul *et al.* 2003). The current study demonstrates that NR2B-NMDARs are responsible for the activation of STEP. This delayed activation of STEP, in turn, can limit the activation of both p38 MAPK (current study) and ERK MAPK (Paul *et al.* 2003). As prolonged activation of either ERK or p38 MAPK has been shown to be involved in neuronal apoptosis (Horstmann *et al.* 1998; Murray *et al.* 1998; Runden *et al.* 1998; Park *et al.* 2002; Stanciu and DeFranco 2002; Poddar and Paul 2009) these findings suggest a role of STEP in mediating the pro-survival effects of NMDA receptor stimulation.

It is not yet known why NR2A-containing NMDARs preferentially activate ERK and p38 MAPK, while STEP is more strongly activated by NR2B-containing receptors. However, one possible mechanism involves differential  $Ca^{2+}$  dynamics triggered by these receptors. NMDA receptor mediated  $Ca^{2+}$  influx is required for activation of both MAPKs and STEP, and we have recently shown that NMDA receptor induced intracellular  $Ca^{2+}$  increases occur in two phases following exposure to high levels of glutamate. An initial rapid but small rise is followed by a gradual but larger

increase. The delayed increase in  $\text{Ca}^{2+}$  is NR2B-NMDAR-dependent and strongly correlates with the delayed activation of STEP (Paul and Connor 2010). The two phase nature of the  $\text{Ca}^{2+}$  increase is also evident from electrophysiological studies in hippocampal neurons demonstrating that exposure to 100  $\mu\text{M}$  NMDA leads to a rapidly developing inward current that would give rise to a rapid  $\text{Ca}^{2+}$  increase, followed by a larger slowly developing inward current that would give rise to a larger more slowly developing  $\text{Ca}^{2+}$  increase (Chen *et al.* 1997; Thompson *et al.* 2008). Taken together, it seems likely that a rapid initial  $\text{Ca}^{2+}$  increase (presumably because of NR2A-NMDA stimulation), is sufficient to activate ERK and p38 MAPK. Higher levels of  $\text{Ca}^{2+}$  influx via NR2B-NMDAR may initially activate cell survival pathways, including the activation of STEP. Consistent with this interpretation several previous studies in cultured neurons demonstrated that moderate increase in intracellular  $\text{Ca}^{2+}$  inhibits cell death induced by oxygen glucose deprivation and serum or growth factor withdrawal (Berridge *et al.* 2000; Chen *et al.* 2003; Cheng *et al.* 2003; Mottet *et al.* 2003; Bickler and Fahlman 2004). Moderate increase in  $\text{Ca}^{2+}$  also appears to be critical in the phenomenon of ischemic preconditioning in which tolerance to a severe ischemic insult is induced by antecedent of mild insult or stimulus (Grabb and Choi 1999; Semenov *et al.* 2002; Raval *et al.* 2003; Tauskela and Morley 2004). The extent to which  $\text{Ca}^{2+}$ -dependent STEP activation contributes to cell survival in these previous studies is not yet known. However, this neuroprotection may become ineffective if the extracellular insult is sustained leading to accumulation of potentially lethal levels of  $\text{Ca}^{2+}$  and activation of detrimental pathways, which down-regulates the pro-survival factors that includes STEP.

A recent study showed that extrasynaptic NMDA receptor stimulation led to calpain-dependent cleavage of STEP, and proposed that this was an important contributor to excitotoxic neuronal injury (Xu *et al.* 2009). The results of our study are consistent with this general conclusion, as proteolytic cleavage of STEP was clearly observed following extended glutamate exposures, and maintaining STEP activity with the TAT-STEP peptide was sufficient to prevent neuronal injury. Furthermore, the TAT-STEP peptide also prevented *in vivo* injury strongly supporting the idea that targeting STEP breakdown could be an effective therapeutic approach.

Our results also reveal an important additional phase of p38 MAPK and STEP activation during glutamate-NMDA receptor stimulation that was not observed in the prior study. Immediately following glutamate treatment, a rapid initial activation of p38 MAPK is observed that is followed by a time-dependent decline because of the interaction of active STEP with p38 MAPK. Earlier studies have also reported activation of STEP following intracellular  $\text{Ca}^{2+}$  increase and activation of calcineurin (Paul *et al.* 2003; Snyder *et al.* 2005). It is difficult to explain why we saw this initial phase of p38 and STEP activation that was not observed in the

study by Xu *et al.* (2009). One possibility is that the stimuli used in the present study (100  $\mu\text{M}$  glutamate) permit more rapid activation of p38 and STEP, compared with the approach of Xu *et al.* (2009) that used 10  $\mu\text{M}$  NMDA along with bicuculline to stimulate extrasynaptic NMDARs.

Based on our data, we suggest that irrespective of its anatomical location, NR2B-NMDAR stimulation can induce dephosphorylation and subsequent activation of STEP. Our hypothesis is that trans-synaptic stimulation of NR2B-NMDAR may lead to activation of STEP very close to the site of  $\text{Ca}^{2+}$  entry that plays a role in modulating synaptic plasticity through down-regulation of ERK and p38 MAPK localized at the synaptic membrane. Whereas an excitotoxic insult leading to a more global increase in  $\text{Ca}^{2+}$  through all NR2B-NMDARs leads to overall increase in STEP activity that promotes neuroprotection by preventing the chronic activation and accumulation of p38 and ERK MAPK in the nucleus. However, an extended insult may lead to STEP's degradation, facilitating cell death pathways.

The regulation of p38 MAPK activity by STEP described in the present study may also contribute to neuronal survival following stroke *in vivo*. A marked increase in extracellular glutamate levels has been observed in severe brain disorders such as ischemic stroke and traumatic brain injury and is responsible for NMDAR induced brain damage (Choi and Rothman 1990; Dirnagl *et al.* 1999). Activation of p38 MAPK has also been observed in the penumbral region following ischemic stroke (Barone *et al.* 2001a,b; Cao *et al.* 2005; Nito *et al.* 2008) and plays a crucial role in ischemic brain injury. It is possible that degradation of active STEP is responsible for p38 MAPK activation in these models. Consistent with this hypothesis, a recent study using an animal model of ischemic stroke has shown that STEP is unilaterally down-regulated in the stroked hemisphere over time. Furthermore, STEP levels are increased in a neuroprotective preconditioning paradigm, in regions of the brain that are resistant to ischemic brain damage (Braithwaite *et al.* 2008). p38 MAPK has been linked to an aberrant increase in expression of cyclooxygenase-2, an enzyme involved in the generation of pro-inflammatory prostanoids and reactive oxygen species (Rockwell *et al.* 2004; Moolwaney and Igwe 2005; Nito *et al.* 2008). Thus, assessing the role of STEP in modulating cyclooxygenase-2 expression and pro-inflammatory responses may be helpful in determining whether the TAT-STEP peptide can be considered as a therapeutic agent to inhibit the progression of brain damage following ischemic stroke or related neurological disorders.

## Acknowledgements

This work was supported by the National Institutes of Health grants NS059962 (Paul, S) and P20 RR015636 (Okada, Y). We would like to thank Dr. William Shuttleworth, University of New Mexico, for his helpful comments and Eduardo Estrada for technical assistance.

## Supporting information

Additional Supporting Information may be found in the online version of this article:

**Figure S1.** Glutamate-induced neuronal cell death is regulated by p38 MAPK.

As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials are peer-reviewed and may be re-organized for online delivery, but are not copy-edited or typeset. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.

## References

- Arundine M. and Tymianski M. (2004) Molecular mechanisms of glutamate-dependent neurodegeneration in ischemia and traumatic brain injury. *Cell. Mol. Life Sci.* **61**, 657–668.
- Barone F. C., Irving E. A., Ray A. M. *et al.* (2001a) Inhibition of p38 mitogen-activated protein kinase provides neuroprotection in cerebral focal ischemia. *Med. Res. Rev.* **21**, 129–145.
- Barone F. C., Irving E. A., Ray A. M. *et al.* (2001b) SB 239063, a second-generation p38 mitogen-activated protein kinase inhibitor, reduces brain injury and neurological deficits in cerebral focal ischemia. *J. Pharmacol. Exp. Ther.* **296**, 312–321.
- Berridge M. J., Lipp P. and Bootman M. D. (2000) The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* **1**, 11–21.
- Bickler P. E. and Fahlman C. S. (2004) Moderate increases in intracellular calcium activate neuroprotective signals in hippocampal neurons. *Neuroscience* **127**, 673–683.
- Bolshakov V. Y., Carboni L., Cobb M. H., Siegelbaum S. A. and Belardetti F. (2000) Dual MAP kinase pathways mediate opposing forms of long-term plasticity at CA3-CA1 synapses. *Nat. Neurosci.* **3**, 1107–1112.
- Boulanger L. M., Lombroso P. J., Raghunathan A., During M. J., Wahle P. and Naegel J. R. (1995) Cellular and molecular characterization of a brain-enriched protein tyrosine phosphatase. *J. Neurosci.* **15**, 1532–1544.
- Braithwaite S. P., Xu J., Leung J., Urfer R., Nikolich K., Oksenberg D., Lombroso P. J. and Shamloo M. (2008) Expression and function of striatal enriched protein tyrosine phosphatase is profoundly altered in cerebral ischemia. *Eur. J. Neurosci.* **27**, 2444–2452.
- Bult A., Zhao F., Dirx Jr. R., Raghunathan A., Solimena M. and Lombroso P. J. (1997) STEP: a family of brain-enriched PTPs. Alternative splicing produces transmembrane, cytosolic and truncated isoforms. *Eur. J. Cell Biol.* **72**, 337–344.
- Cao J., Viholainen J. I., Dart C., Warwick H. K., Leyland M. L. and Courtney M. J. (2005) The PSD95-nNOS interface: a target for inhibition of excitotoxic p38 stress-activated protein kinase activation and cell death. *J. Cell Biol.* **168**, 117–126.
- Chen Q. X., Perkins K. L., Choi D. W. and Wong R. K. (1997) Secondary activation of a cation conductance is responsible for NMDA toxicity in acutely isolated hippocampal neurons. *J. Neurosci.* **17**, 4032–4036.
- Chen W. G., Chang Q., Lin Y., Meissner A., West A. E., Griffith E. C., Jaenisch R. and Greenberg M. E. (2003) Derepression of BDNF transcription involves calcium-dependent phosphorylation of MeCP2. *Science (New York, NY)* **302**, 885–889.
- Cheng A., Wang S., Yang D., Xiao R. and Mattson M. P. (2003) Calmodulin mediates brain-derived neurotrophic factor cell survival signaling upstream of Akt kinase in embryonic neocortical neurons. *J. Biol. Chem.* **278**, 7591–7599.
- Choi D. W. and Rothman S. M. (1990) The role of glutamate neurotoxicity in hypoxic-ischemic neuronal death. *Annu. Rev. Neurosci.* **13**, 171–182.
- Dirnagl U., Iadecola C. and Moskowitz M. A. (1999) Pathobiology of ischaemic stroke: an integrated view. *Trends Neurosci.* **22**, 391–397.
- Gisabella B., Rowan M. J. and Anwyl R. (2003) Mechanisms underlying the inhibition of long-term potentiation by preconditioning stimulation in the hippocampus in vitro. *Neuroscience* **121**, 297–305.
- Grabb M. C. and Choi D. W. (1999) Ischemic tolerance in murine cortical cell culture: critical role for NMDA receptors. *J. Neurosci.* **19**, 1657–1662.
- Haelewyn B., Alix P., Maubert E. and Abraini J. H. (2008) NMDA-induced striatal brain damage and time-dependence reliability of thionin staining in rats. *J. Neurosci. Methods* **168**, 479–482.
- Horstmann S., Kahle P. J. and Borasio G. D. (1998) Inhibitors of p38 mitogen-activated protein kinase promote neuronal survival in vitro. *J. Neurosci. Res.* **52**, 483–490.
- Iadecola C., Niwa K., Nogawa S., Zhao X., Nagayama M., Araki E., Morham S. and Ross M. E. (2001) Reduced susceptibility to ischemic brain injury and N-methyl-D-aspartate-mediated neurotoxicity in cyclooxygenase-2-deficient mice. *Proc. Natl Acad. Sci. USA* **98**, 1294–1299.
- Ingram A. J., James L., Thai K., Ly H., Cai L. and Scholey J. W. (2000) Nitric oxide modulates mechanical strain-induced activation of p38 MAPK in mesangial cells. *Am. J. Physiol. Renal Physiol.* **279**, F243–F251.
- Kawasaki H., Morooka T., Shimohama S., Kimura J., Hirano T., Gotoh Y. and Nishida E. (1997) Activation and involvement of p38 mitogen-activated protein kinase in glutamate-induced apoptosis in rat cerebellar granule cells. *J. Biol. Chem.* **272**, 18518–18521.
- Kim M. J., Dunah A. W., Wang Y. T. and Sheng M. (2005) Differential roles of NR2A- and NR2B-containing NMDA receptors in Ras-ERK signaling and AMPA receptor trafficking. *Neuron* **46**, 745–760.
- Kohr G. (2006) NMDA receptor function: subunit composition versus spatial distribution. *Cell Tissue Res.* **326**, 439–446.
- Kumar S., Boehm J. and Lee J. C. (2003) p38 MAP kinases: key signalling molecules as therapeutic targets for inflammatory diseases. *Nat. Rev. Drug Discov.* **2**, 717–726.
- Lipton S. A. and Rosenberg P. A. (1994) Excitatory amino acids as a final common pathway for neurologic disorders. *N. Eng. J. Med.* **330**, 613–622.
- Liu Y., Wong T. P., Aarts M. *et al.* (2007) NMDA receptor subunits have differential roles in mediating excitotoxic neuronal death both in vitro and in vivo. *J. Neurosci.* **27**, 2846–2857.
- Luo J., Wang Y., Yasuda R. P., Dunah A. W. and Wolfe B. B. (1997) The majority of N-methyl-D-aspartate receptor complexes in adult rat cerebral cortex contain at least three different subunits (NR1/NR2A/NR2B). *Mol. Pharmacol.* **51**, 79–86.
- Moolwaney A. S. and Igwe O. J. (2005) Regulation of the cyclooxygenase-2 system by interleukin-1beta through mitogen-activated protein kinase signaling pathways: a comparative study of human neuroglioma and neuroblastoma cells. *Brain Res. Mol. Brain Res.* **137**, 202–212.
- Mottet D., Michel G., Renard P., Ninane N., Raes M. and Michiels C. (2003) Role of ERK and calcium in the hypoxia-induced activation of HIF-1. *J. Cell. Physiol.* **194**, 30–44.
- Mukherjee P. K., DeCoster M. A., Campbell F. Z., Davis R. J. and Bazan N. G. (1999) Glutamate receptor signaling interplay modulates stress-sensitive mitogen-activated protein kinases and neuronal cell death. *J. Biol. Chem.* **274**, 6493–6498.
- Murray B., Alessandrini A., Cole A. J., Yee A. G. and Furshpan E. J. (1998) Inhibition of the p44/42 MAP kinase pathway protects

- hippocampal neurons in a cell-culture model of seizure activity. *Proc. Natl Acad. Sci. USA* **95**, 11975–11980.
- Nguyen T. H., Liu J. and Lombroso P. J. (2002) Striatal enriched phosphatase 61 dephosphorylates Fyn at phosphotyrosine 420. *J. Biol. Chem.* **277**, 24274–24279.
- Nito C., Kamada H., Endo H., Niizuma K., Myer D. J. and Chan P. H. (2008) Role of the p38 mitogen-activated protein kinase/cytosolic phospholipase A2 signaling pathway in blood-brain barrier disruption after focal cerebral ischemia and reperfusion. *J. Cereb. Blood Flow Metab.* **28**, 1686–1696.
- Park J. G., Yuk Y., Rhim H., Yi S. Y. and Yoo Y. S. (2002) Role of p38 MAPK in the regulation of apoptosis signaling induced by TNF- $\alpha$  in differentiated PC12 cells. *J. Biochem. Mol. Biol.* **35**, 267–272.
- Paul S. and Connor J. (2010) NR2B-NMDA receptor mediated increases in intracellular Ca<sup>2+</sup> concentration regulate the tyrosine phosphatase, STEP, and ERK MAP kinase signaling. *J. Neurochem.* **114**, 1107–1118.
- Paul S., Snyder G. L., Yokakura H., Picciotto M. R., Nairn A. C. and Lombroso P. J. (2000) The Dopamine/D1 receptor mediates the phosphorylation and inactivation of the protein tyrosine phosphatase STEP via a PKA-dependent pathway. *J. Neurosci.* **20**, 5630–5638.
- Paul S., Nairn A. C., Wang P. and Lombroso P. J. (2003) NMDA-mediated activation of the tyrosine phosphatase STEP regulates the duration of ERK signaling. *Nat. Neurosci.* **6**, 34–42.
- Paul S., Olausson P., Venkataramani D. V. *et al.* (2007) The striatal-enriched protein tyrosine phosphatase gates long-term potentiation and fear memory in the lateral amygdala. *Biol. Psychiatry* **61**, 1049–1061.
- Paxinos G. and Watson C. (1986). *The Rat Brain in Stereotaxic Coordinates*. Academic Press, New York.
- Poddar R. and Paul S. (2009) Homocysteine-NMDA receptor-mediated activation of extracellular signal-regulated kinase leads to neuronal cell death. *J. Neurochem.* **110**, 1095–1106.
- Pulido R., Zuniga A. and Ullrich A. (1998) PTP-SL and STEP protein tyrosine phosphatases regulate the activation of the extracellular signal-regulated kinases ERK1 and ERK2 by association through a kinase interaction motif. *EMBO J.* **17**, 7337–7350.
- Raval A. P., Dave K. R., Mochly-Rosen D., Sick T. J. and Perez-Pinzon M. A. (2003) Epsilon PKC is required for the induction of tolerance by ischemic and NMDA-mediated preconditioning in the organotypic hippocampal slice. *J. Neurosci.* **23**, 384–391.
- Rockwell P., Martinez J., Papa L. and Gomes E. (2004) Redox regulates COX-2 upregulation and cell death in the neuronal response to cadmium. *Cell. Signal.* **16**, 343–353.
- Rodrigues S. M., Schafe G. E. and LeDoux J. E. (2001) Intra-amygdala blockade of the NR2B subunit of the NMDA receptor disrupts the acquisition but not the expression of fear conditioning. *J. Neurosci.* **21**, 6889–6896.
- Runden E., Seglen P. O., Haug F. M., Ottersen O. P., Wieloch T., Shamloo M. and Laake J. H. (1998) Regional selective neuronal degeneration after protein phosphatase inhibition in hippocampal slice cultures: evidence for a MAP kinase-dependent mechanism. *J. Neurosci.* **18**, 7296–7305.
- Semenov D. G., Samoilov M. O. and Lazarewicz J. W. (2002) Calcium transients in the model of rapidly induced anoxic tolerance in rat cortical slices: involvement of NMDA receptors. *Neuro-Signals* **11**, 329–335.
- Snyder E. M., Nong Y., Almeida C. G. *et al.* (2005) Regulation of NMDA receptor trafficking by amyloid-beta. *Nat. Neurosci.* **8**, 1051–1058.
- Stanciu M. and DeFranco D. B. (2002) Prolonged nuclear retention of activated extracellular signal-regulated protein kinase promotes cell death generated by oxidative toxicity or proteasome inhibition in a neuronal cell line. *J. Biol. Chem.* **277**, 4010–4017.
- Tauskela J. S. and Morley P. (2004) On the role of Ca<sup>2+</sup> in cerebral ischemic preconditioning. *Cell Calcium* **36**, 313–322.
- Thomas G. M. and Huganir R. L. (2004) MAPK cascade signalling and synaptic plasticity. *Nat. Rev. Neurosci.* **5**, 173–183.
- Thompson R. J., Jackson M. F., Olah M. E., Rungta R. L., Hines D. J., Beazely M. A., MacDonald J. F. and MacVicar B. A. (2008) Activation of pannexin-1 hemichannels augments aberrant bursting in the hippocampus. *Science (New York, NY)* **322**, 1555–1559.
- Tovar K. R. and Westbrook G. L. (1999) The incorporation of NMDA receptors with a distinct subunit composition at nascent hippocampal synapses in vitro. *J. Neurosci.* **19**, 4180–4188.
- Tu W., Xu X., Peng L. *et al.* (2010) DAPK1 interaction with NMDA receptor NR2B subunits mediates brain damage in stroke. *Cell* **140**, 222–234.
- Walker D. L. and Davis M. (2008) Amygdala infusions of an NR2B-selective or an NR2A-preferring NMDA receptor antagonist differentially influence fear conditioning and expression in the fear-potentiated startle test. *Learn. Mem. (Cold Spring Harbor, NY)* **15**, 67–74.
- Wang Q., Walsh D. M., Rowan M. J., Selkoe D. J. and Anwyl R. (2004) Block of long-term potentiation by naturally secreted and synthetic amyloid beta-peptide in hippocampal slices is mediated via activation of the kinases c-Jun N-terminal kinase, cyclin-dependent kinase 5, and p38 mitogen-activated protein kinase as well as metabotropic glutamate receptor type 5. *J. Neurosci.* **24**, 3370–3378.
- Waxman E. A. and Lynch D. R. (2005) N-methyl-D-aspartate receptor subtype mediated bidirectional control of p38 mitogen-activated protein kinase. *J. Biol. Chem.* **280**, 29322–29333.
- Xu J., Kurup P., Zhang Y., Goebel-Goody S. M., Wu P. H., Hawasli A. H., Baum M. L., Bibb J. A. and Lombroso P. J. (2009) Extrasynaptic NMDA receptors couple preferentially to excitotoxicity via calpain-mediated cleavage of STEP. *J. Neurosci.* **29**, 9330–9343.
- Zhao M. G., Toyoda H., Lee Y. S. *et al.* (2005) Roles of NMDA NR2B subtype receptor in pre long-term potentiation and contextual fear memory. *Neuron* **47**, 859–872.
- Zhu J. J., Qin Y., Zhao M., Van Aelst L. and Malinow R. (2002) Ras and Rap control AMPA receptor trafficking during synaptic plasticity. *Cell* **110**, 443–455.