

Efficient Test to Demonstrate Genuine Three Particle Nonlocality

Kaushiki Mukherjee,^{1,*} Biswajit Paul,^{2,†} and Debasis Sarkar^{1,‡}

¹*Department of Applied Mathematics, University of Calcutta, 92, A.P.C. Road, Kolkata-700009, India.*

²*Department of Mathematics, St.Thomas' College of Engineering and Technology,
4, Diamond Harbour Road, Alipore, Kolkata-700023, India.*

(Dated: October 8, 2018)

According to the studies of genuine tripartite nonlocality in discrete variable quantum systems conducted so far, Svetlichny inequality is considered as the best Bell-type inequality to detect genuine (three way) nonlocality of pure tripartite genuine entangled states. In the present work, we have considered another Bell-type inequality (which has been reported as the 99-th facet of NS_2 local polytope in (J.-D. Bancal, et.al., Phys. Rev.A **88**, 014102 (2013)), to reveal genuine tripartite nonlocality of generalized GHZ (Greenberger-Horne-Zeilinger) class and a subclass of extended GHZ class states([1]) thereby proving the conjecture given by Bancal, et.al.[31] for the GGHZ class and the subclass of extended GHZ states. We compare the violation of this inequality with Svetlichny inequality which reveals the efficiency of the former inequality over the latter to demonstrate genuine nonlocality using the above classes of quantum states. Even in some cases discord monogamy score can be used as a better measure of quantum correlation over Svetlichny inequality for those classes of pure states. Besides, the 99-th facet inequality is found efficient not only for revealing genuine nonlocal behavior of correlations emerging in systems using pure entangled states but also in some cases of mixed entangled states over Svetlichny inequality and some well known measures of entanglement .

PACS numbers: 03.65.Ud, 03.67.-a

I. MOTIVATION

The correlated statistics arising by performing local measurements on an entangled state are nonlocal in nature in the sense that they violate a Bell inequality [2, 3]. Nonlocality has been a core part of quantum mechanics which has been supported by many experimental evidences[4, 5]. So far there has been a lot of analysis about the nonlocal nature of correlations arising between two space-like separated quantum systems exploring different tools demonstrating nonlocality in bipartite systems, such as, different forms of Bell-type inequalities, nonlocality witnesses, etc. However, the understanding about multipartite nonlocality has not yet reached a satisfactory level due to increasing complexity while one shift from a bipartite to a multipartite scenario. In this context, perhaps the most interesting topic is that of genuine nonlocality. It may be referred to as correlations emerging in a n -party quantum system when all of the spatially separated parties constituting the sys-

tem are nonlocally correlated. It is the strongest form of nonlocality. Apart from enriching quantum mechanics theoretically, study of nonlocality has significant contributions in the field of applications, for example, developing practical quantum information processing protocols [6–15], communication complexity problems[16], device independent quantum cryptography [17–20], randomness expansion [21, 22], measurement-based quantum computation [23, 24], etc. Besides, multipartite long-range correlations can play an important role in various condensed matter systems[25, 26], quantum phase transitions [11, 27, 28, 72]. Svetlichny laid the cornerstone in the study of genuine multipartite nonlocality.

A. Svetlichny inequality

Svetlichny introduced a different form of correlations namely hybrid local-nonlocal form:

$$P(abc/XYZ) = \sum_{\lambda} q_{\lambda} P_{\lambda}(ab/XY) P_{\lambda}(c/Z) + \sum_{\mu} q_{\mu} P_{\mu}(ac/XZ) P_{\mu}(b/Y) + \sum_{\nu} q_{\nu} P_{\nu}(bc/YZ) P_{\nu}(a/X); \quad (1)$$

where $a, b, c \in \{0, 1\}$ denote the outputs and $X, Y, Z \in \{0, 1\}$ denote the inputs of the three parties Alice, Bob and Charlie respectively. Here $0 \leq q_{\lambda}, q_{\mu}, q_{\nu} \leq 1$ and $\sum_{\lambda} q_{\lambda} + \sum_{\mu} q_{\mu} + \sum_{\nu} q_{\nu} = 1$. The correlations which can-

not be written in this form are genuine nonlocal (we will refer it as, S_2 (Svetlichny) nonlocal, in future). In order to detect genuine nonlocality, a Bell-type inequality $S \leq 4$ is provided where

$$S = \langle X_0 Y_0 Z_0 \rangle + \langle X_1 Y_0 Z_0 \rangle - \langle X_0 Y_1 Z_0 \rangle + \langle X_1 Y_1 Z_0 \rangle + \langle X_0 Y_0 Z_1 \rangle - \langle X_1 Y_0 Z_1 \rangle + \langle X_0 Y_1 Z_1 \rangle + \langle X_1 Y_1 Z_1 \rangle, \quad (2)$$

violation of which guarantees *Svetlichny* nonlocality ([30]) (S_2 nonlocality ([31])). Svetlichny inequality has been further generalized to arbitrary number of parties [32, 33] and arbitrary dimension [34, 35]. This inequality (2) is violated by GHZ and W states [36–38] and can be regarded as a unique witness of genuine tripartite nonlocality in a scenario where each of the three parties performs two dichotomic measurements [30]. However, it can be regarded as a sufficient criteria only for detecting genuine nonlocality as there exist some genuine nonlocal correlations which satisfy this inequality [31, 39, 40].

B. Incompleteness of S_2 correlations

The definition (1) being general, no restrictions were imposed on the bipartite terms used in the inequality

(2). There may be one-way or both way signaling between a pair of parties or both the parties may perform simultaneous measurements. Thus, Svetlichny type nonlocality and hence the inequality (2) lacks physical motivation which in turn may lead to grandfather-style paradoxes [31] and inconsistency in operational purposes [39, 40]. In [31] Bancal et.al., removed this sort of ambiguity by putting restrictions on the bipartite terms and thereby provided two alternative definitions of nonlocality: NS_2 (no signaling) type and T_2 (time ordering) type.

C. NS_2 correlations and Bancal et.al.'s conjecture

Among these two types, in the NS_2 type nonlocality, bipartite correlation terms abide by the no signaling criteria:

$$P(abc/XYZ) = \sum_{\lambda} q_{\lambda} P_{\lambda}(ab/XY) P_{\lambda}(c/Z) + \sum_{\mu} q_{\mu} P_{\mu}(ac/XZ) P_{\mu}(b/Y) + \sum_{\nu} q_{\nu} P_{\nu}(bc/YZ) P_{\nu}(a/X) \quad (3)$$

where the bipartite terms satisfy no signaling conditions of the form:

$$P_{\lambda}(a/XY) = P_{\lambda}(a/XY') \quad \forall a, X, Y, Y' \quad (4)$$

$$P_{\lambda}(b/XY) = P_{\lambda}(b/X'Y) \quad \forall b, X, X', Y. \quad (5)$$

The correlations which are of the above form are called NS_2 (no signaling) local. Otherwise, they are NS_2 nonlocal. The no signaling constraints imposed on the bipartite terms exclude the possibility of the outcomes of one or two parties influencing the inputs of the remaining one. Hence this form of nonlocality is in general weaker than that of Svetlichny nonlocality. In [31], 185 Bell-type inequalities are given which constitute the full class of facets of NS_2 local polytope. Violation of any of these facets (Bell-type inequalities) guarantees NS_2 nonlocality. Svetlichny inequality constitute the 185-th class. So far, mostly the 185-th class facet inequality, i.e., Svetlichny inequality (2) has been used as a tool to demonstrate genuine nonlocality. For instance, in [36] S. Ghose et.al., analyzed the relation between genuine nonlocality and three-tangle (genuine tripartite entanglement measure) [41] for the class of GGHZ [1, 42] and Maximal Slice (MS) states [43]. The latter are included in the subclass S of extended GHZ states [1, 42]). They showed

that up to projective measurements, GGHZ states show genuine nonlocality for three-tangle (τ) greater than $\frac{1}{3}$ whereas MS states violate Svetlichny inequality for any positive value of τ . In [37], closed forms of the bounds of Svetlichny inequality were derived for extended GHZ and W class of states. In the framework where a complete set of Bell-type inequalities were introduced by Bancal et. al. ([31]), it was conjectured that genuine tripartite entanglement of a pure state guarantees genuine nonlocality. With the help of one of these Bell-type inequalities (168-th class), Adesso et. al., investigated the genuine tripartite nonlocality of three-mode Gaussian states in continuous variable systems [44]. Besides, in [45] Almeida et.al., provided sufficient criteria for a quantum system to be fully nonlocal according to a given partition and also to be genuinely multipartite nonlocal. Other than the statistical approach [30, 31] genuine nonlocality of tripartite states was revealed with the aid of Hardy type argument [46–48]. Apart from demonstrating nonlocality in quantum systems, an information theoretic measure of discord monogamy score [49, 50] has been used to exploit quantum correlations beyond entanglement for multipartite systems. Although all these studies have contributed in providing different means of demonstrating nonlocality and thereby quantumness in different physical systems, yet, it is still a matter of interest to develop better tools

for exploiting the same. Our work basically focuses in this direction.

D. Summary of the work

We have used one of the facets of NS_2 local polytope as a better tool than Svetlichny for demonstrating three way nonlocal nature of correlations emerging in discrete variable quantum systems. This in turn is useful to analyze the existing relation in between nonlocal nature of the resulting correlations with that of genuine tripartite entanglement of the quantum states involved in the systems. We have used the 99-th class facet of the NS_2 local polytope [31]. The 99-th facet inequality ([31]) is given by:

$$NS \leq 3. \quad (6)$$

where $NS = \langle X_1 Y_1 \rangle + \langle X_0 Y_0 Z_0 \rangle + \langle Y_1 Z_0 \rangle + \langle X_1 Z_1 \rangle - \langle X_0 Y_0 Z_1 \rangle$. For this facet, we derive the closed form of maximum violation for two class of pure tripartite states: GHZ and a subclass (S) of extended GHZ states ([1]) under projective measurements. Interestingly the 99-th class facet (Eq.6) helps us to reveal genuine nonlocality of GHZ states for any non zero value of τ unlike Eq(2), where the same is revealed only for $\tau > \frac{1}{3}$. This in turn proves the conjecture made by Bancal et.al.,([31]) for the GHZ class of states. Further, a comparative study of the violations of 99-th facet inequality and Svetlichny inequality for the subclass S of extended GHZ states reveals that the former(6) gives advantage over the latter for a certain range of τ . In particular, MS class of states (a subclass of S) exhibits genuine tripartite nonlocality for any positive amount of τ for both of these facet inequalities. Consequently, 99-th class facet inequality(6) emerges as a more efficient tool compared to Svetlichny for revealing three-way nonlocality in a quantum system using the above mentioned two classes of pure tripartite states. Quantumness is revealed in a system via generation of nonlocal correlations. From that perspective 99-th facet inequality serves as a good measure of quantum correlation behaving in a similar manner as discord monogamy score (δ_D) for the complete GHZ class and also for subclass S of extended GHZ class. The similarity in the pattern of variation of discord monogamy score (δ_D) and that of amount of violation of 99-th facet inequality with the state parameter or amount of entanglement also supports the fact that δ_D helps us to detect genuine nonlocality via revelation of quantumness even in some cases where Svetlichny inequality cannot be used as a tool to detect genuine nonlocality of these two classes of pure states. For instance, considering GHZ class of states, $\delta_D > 0$ in the subinterval $[0, 0.393]$ of the interval $[0, \frac{\pi}{4}]$ of the state parameter (η) where the class reveals nonlocality (guaranteed by violation of the 99-th facet inequality) but Svetlichny cannot be used to demonstrate the same. Similar sort of results also exist for subclass(S)

of extended GHZ class of states. In case of mixed tripartite states, the main focus so far was on analyzing the entanglement properties of the states via various entanglement measures. Here we have studied nonlocal behavior of noisy GHZ states, noisy MS states and some family of high rank mixed states(whose explicit expressions for the three-tangle(τ) are reported in [55–59]) by providing closed forms of violation of 99th facet inequality(6). Our work can be organized as follows: Section (II) deals with some mathematical prerequisites. Bounds for violation of the 99th facet(6) are discussed in Section(III). Next three sections(IV-VI) deal with various sides of getting advantage of the 99th facet over Svetlichny inequality(2) from physical point of view. In Section(VII) we end up with a conclusion.

II. PRELIMINARIES

To begin with we first explain the three-tangle (τ) which is used as the measure of genuine tripartite entanglement [41]:

$$\tau = \mathcal{C}_{1(23)}^2 - \mathcal{C}_{12}^2 - \mathcal{C}_{13}^2 \quad (7)$$

$\mathcal{C}_{1(23)}^2$ is a measure of entanglement between the first qubit(1) and the joint state of the last two qubits (2, 3). \mathcal{C}_{12}^2 and \mathcal{C}_{13}^2 are the concurrences [60] measuring nature of bipartite entanglement between qubits 1, 2 and 1, 3 respectively. τ remains invariant under permutation of the indices (1, 2, 3) and lies in the interval $[0, 1]$. For separable and bi-separable states $\tau = 0$ and for GHZ state, it is 1. For a three-qubit pure state of the form:

$$|\psi\rangle = a|011\rangle + b|101\rangle + c|110\rangle + d|000\rangle + he^{i\gamma}|111\rangle \quad (8)$$

three-tangle is given by

$$\tau = 4d\sqrt{(dh^2 - 4abc)^2 + 16abcdh^2\cos^2\gamma} \quad (9)$$

Quantum Monogamy Score: Monogamy constrains the sharing of quantum correlations among subsystems of a multipartite quantum state. For a tripartite quantum state ρ_{ABC} , a bipartite quantum correlation measure (\mathcal{Q}) is monogamous (with A as the “nodal observer”) if

$$\mathcal{Q}(\rho_{A:BC}) \geq \mathcal{Q}(\rho_{AB}) + \mathcal{Q}(\rho_{AC}). \quad (10)$$

Here, $\mathcal{Q}(\rho_{AB})$, $\mathcal{Q}(\rho_{AC})$ and $\mathcal{Q}(\rho_{A:BC})$ denotes quantum correlation (considering \mathcal{Q} as a bipartite measure) between subsystems (A, B), (A, C) and between system A and subsystems B and C taken together respectively. The above inequality can be recast as

$$\delta_{\mathcal{Q}} \equiv \mathcal{Q}(\rho_{A:BC}) - \mathcal{Q}(\rho_{AB}) - \mathcal{Q}(\rho_{AC}) \geq 0. \quad (11)$$

$\delta_{\mathcal{Q}}$ is regarded as the monogamy score. If discord (D) is considered as the quantum correlation measure (\mathcal{Q}) then we get discord monogamy score [49, 50]:

$$\delta_D = D(\rho_{A:BC}) - D(\rho_{AB}) - D(\rho_{AC}) \quad (12)$$

where $D(\rho_{AB})$ denotes quantum discord between subsystems A and B [61–63]. Discord monogamy score (δ_D) has been interpreted as a multipartite information-theoretic quantum correlation measure [50]. A tripartite state ρ_{ABC} satisfies monogamy relation (Eq.(10)), if $\delta_D > 0$, whereas it violates Eq.10, if $\delta_D \leq 0$.

III. BOUNDS FOR BELL INEQUALITY VIOLATION

A. GGHZ class

First we consider generalized Greenberger-Horne-Zeilinger (GGHZ) class of states [1]:

$$|\varphi_{GGHZ}\rangle = \cos\eta|000\rangle + \sin\eta|111\rangle \text{ where } \eta \in [0, \frac{\pi}{4}]. \quad (13)$$

The GHZ states([64]) are for $\eta = \frac{\pi}{4}$. The GGHZ states are significant for the study of nonlocality as they are the only class of pure three qubit states such that all information about them can be coded in the corresponding tripartite correlations (they cannot be uniquely reconstructed from the corresponding reduced two-qubit states) [65]. Besides, because of inherent symmetry in their structure [43], they possess some unique entanglement characteristics for which they are useful in various information processing protocols. For this class of states the bound of Eq.(6) is given by(see Appendix.A),

$$S \leq B_1 \text{ where } B_1 = 1 + 2\sqrt{1 + \sin^2 2\eta}. \quad (14)$$

Here $\tau = \sin^2 2\eta$. Except for $\eta = 0$, the class of states (13) has genuine tripartite entanglement. Thus, from Eq.(14) we have:

$$B_1 = 1 + 2\sqrt{1 + \tau}. \quad (15)$$

The closed form of violation of Svetlichny inequality is given by ([36]) $S < B_2$ where

$$\begin{aligned} B_2 &= 4\sqrt{1 - \tau} \quad \text{if } \tau \leq \frac{1}{3} \\ &= 4\sqrt{2\tau} \quad \text{if } \tau \geq \frac{1}{3}. \end{aligned} \quad (16)$$

Hence for any amount of genuine tripartite entanglement ($\tau > 0$), genuine nonlocality is revealed (using 99-th facet inequality), i.e., $B_1 > 3$ (FIG.2). Whereas the restrictions imposed on the amount of tripartite entanglement is $\tau \in (\frac{1}{3}, 1]$ to show genuine nonlocality (FIG.1) when Svetlichny inequality is used [36].




FIG. 1: The variation of violation of Svetlichny inequality with three-tangle τ is plotted for GGHZ (dashed line) and MS (solid line) subclasses ([36]) of states. MS violates Svetlichny for any value of τ unlike that of GGHZ class which reveals genuine nonlocality (S_2 nonlocality) for $\tau > \frac{1}{3}$.




FIG. 2: The amount of nonlocality in terms of the 99-th class facet inequality violation (N_{S_2} -nonlocality ([31]) increases monotonically with the amount of tripartite entanglement (τ) for both GGHZ and MS subclasses. This figure shows that violation is obtained for any amount of three-tangle.

1. Bell inequality violation as a measure of quantumness

It has already been mentioned that as violation of a Bell inequality guarantees nonlocal behaviour of a physical system, hence it may be considered as a mean of detecting quantumness(nonclassicality) of a quantum state. So from that perspective violation of a Bell inequality may be compared with a standard measure of quantumness. For that purpose we have considered here discord monogamy score. For GGHZ class, discord monogamy score(see Appendix.C) is given by:

$$\delta_D = -(\cos^2 \eta \log_2(\cos^2 \eta) + \sin^2 \eta \log_2(\sin^2 \eta)) \quad (17)$$

Both B_1 and δ_D are functions of state parameter η .

Figure 3 shows that the curve of the quantum correlation measure (δ_D) and that of the bound (B_1) both vary in a similar fashion. For $\eta \in [0, \frac{\pi}{4}]$ Svetlichny inequality (2) is not violated in the sub interval $[0, 0.393]$ whereas states show nonlocal character by violating 99-th facet inequality (6). Also, within this sub interval the states have positive discord monogamy score (δ_D). The states having higher measure of quantum correlation show greater violation of Eq.(6) unlike that of violation of Svetlichny inequality (2) (FIG.3).




FIG. 3: The degree of violation of 99-th facet inequality (6), Svetlichny inequality (2) and discord monogamy score (δ_D) are plotted against state parameter η . Clearly, the nature of the curve representing degree of violation (6) is the same as that of δ_D (17) curve which in turn implies that the degree of violation of Eq.(6) is more for the states having greater value of δ_D . However, if violation of Eq.(2) is considered, there exist states belonging to the GGHZ class for which $\delta_D > 0$ but those states cannot reveal genuine nonlocality in terms of violation of Svetlichny inequality (2).

B. Extended GHZ class

The subclass (S) of extended GHZ class of states has the form([1]):

$$|\chi\rangle = \lambda_0|000\rangle + \lambda_3|110\rangle + \lambda_4|111\rangle \quad (18)$$

where, $\lambda_i \in [-1, 1]$, and $\sum_{i=0,3,4} \lambda_i^2 = 1$.

For $\lambda_0 = \frac{1}{\sqrt{2}}$, $\lambda_3 = \frac{\cos \eta}{\sqrt{2}}$ and $\lambda_4 = \frac{\sin \eta}{\sqrt{2}}$, we get the MS states([43]):

$$\frac{1}{\sqrt{2}}(|000\rangle + \cos \eta|110\rangle + \sin \eta|111\rangle) \quad \eta \in [0, \frac{\pi}{4}]. \quad (19)$$

1. Bounds for MS subclass

The MS subclass of extended GHZ states have various practical applications. For instance the maximally

entangled GHZ state: $\frac{|000\rangle + |111\rangle}{\sqrt{2}}$ (belonging to both MS and GGHZ subclasses) has been used in various physical processes [13–15, 66]. For MS state, the NS bound is given by (see Appendix-B):

$$B_3 = 1 + 2\sqrt{1 + \sin^2 \eta}. \quad (20)$$

Here, $\tau = \sin^2 \eta$. Using this relation, the above bound gets modified as:

$$B_3 = 1 + 2\sqrt{1 + \tau}.$$

Hence B_3 is of the same form as that of B_1 (15). Another class of MS states obtained by swapping qubits of second and third parties yield the same bound (B_3). So any amount of genuine tripartite entanglement suffices to produce violation of Eq.(6). A comparison of the different values of the bounds (B_1 , B_3) obtained by using GGHZ and MS class suggest that for $\sin^2 \eta > 0.75$ GGHZ class yields more nonlocality in terms of violation of Eq.(6). Besides, for a given amount of genuine tripartite entanglement both the classes give the same amount of violation of Eq.(6) in contrast to the violation of Svetlichny inequality where MS gives more violation than GGHZ [36, 67, 68]. It is also interesting to note that variation of maximum quantum violation of 99-th facet inequality (upto projective measurements) by a tripartite state belonging to GGHZ or MS class with that of tripartite entanglement measure τ is analogous to variation of maximum quantum violation of CHSH inequality [69] ($2\sqrt{1 + C_{12}^2}$) for two-qubit pure states [70, 71] with that of bipartite entanglement measure C_{12}^2 (FIG.4).




FIG. 4: The red curve gives variation of maximum quantum violation(T) of 99-th facet inequality (6) with that of τ whereas the dashed curve represents variation of maximum quantum violation(B) of CHSH inequality with C_{12}^2 . Evidently, both the violations vary with the corresponding entanglement measures in a similar fashion.

2. Bounds for subclass S

The amount of tripartite entanglement of the subclass S of extended GHZ states (18) is: $\tau = 4\lambda_0^2\lambda_4^2$ and the

measure of bipartite entanglements are: $C_{12}^2 = 4\lambda_3^2\lambda_0^2$, $C_{13}^2 = 0$ and $C_{23}^2 = 0$. The bound for Svetlichny inequality is ([37]):

$$S \leq B_4$$

where

$$\begin{aligned} B_4 &= 4\sqrt{1-\tau}, \quad \text{if } \tau \leq \frac{1-C_{12}^2}{3} \\ &= 4\sqrt{C_{12}^2+2\tau}, \quad \text{if } \tau > \frac{1-C_{12}^2}{3}. \end{aligned} \quad (21)$$

and for the 99-th facet inequality (6) the bound (see Appendix-B) is:

$$S \leq B_5$$

where

$$\begin{aligned} B_5 &= 3 \quad \text{if } \tau \leq \frac{C_{12}^2(1-C_{12}^2)}{1+C_{12}^2}, \\ &= 1 + \sqrt{A+2C} + \sqrt{A-2C} \quad \text{if } \tau > \frac{C_{12}^2(1-C_{12}^2)}{1+C_{12}^2} \end{aligned} \quad (22)$$

where, $A = 1 + \tau$, $C = \sqrt{C_{12}^2(1-\tau-C_{12}^2)}$. As, $4\sqrt{1-\tau} \leq 4$, this class violates Svetlichny inequality (2) if $C_{12}^2 + 2\tau > 1$. Hence up to projective measurements, S_2 nonlocality is revealed in a quantum system using this subclass (S) only if $\tau \in (\frac{1-C_{12}^2}{2}, 1]$. But for the states belonging to S for which $\lambda_0 = \frac{1}{\sqrt{2}}$, S_2 nonlocality is obtained for any positive value of τ (See Appendix D). As the Maximal Slice states (19) belongs to this category, the results related to nonlocality of MS states reported in [36] can be easily recovered from here. Now, 99-th facet inequality is violated if $B_5 > 3$, which in turn implies that τ must lie in $(\frac{C_{12}^2(1-C_{12}^2)}{1+C_{12}^2}, 1]$. Clearly any tripartite state belonging to the subclass S and characterized by $\lambda_0 = \frac{1}{\sqrt{2}}$ (See Appendix D) violates 99-th facet inequality(6). But in general for any state belonging to the subclass S , unlike Svetlichny inequality (2), the 99-th facet inequality (6) helps us to reveal NS_2 nonlocality of the correlations emerging from the quantum systems using the states of the subclass S of extended GHZ class whose tripartite entanglement is characterized by τ lying in the interval $(\frac{C_{12}^2(1-C_{12}^2)}{1+C_{12}^2}, \frac{1-C_{12}^2}{2}]$ (FIG.5). Clearly,

for any amount of bipartite entanglement C_{12}^2 there is a region of advantage of 99-th facet inequality (6) over Svetlichny inequality (2).

3. Measure of quantumness

The discord monogamy score (δ_D) of $|\chi\rangle$ (see Appendix C) is:

$$\frac{((1-\sqrt{1-\tau})\ln(\frac{1-\sqrt{1-\tau}}{2}) + (1+\sqrt{1-\tau})\ln(\frac{1+\sqrt{1-\tau}}{2}))}{\ln(4)}. \quad (23)$$




FIG. 5: The shaded region indicates the area where 99-th facet inequality (6) emerges as a more efficient tool over Svetlichny inequality (2) for revealing nonlocal nature of tripartite correlations in a quantum system using any state of the subclass S of the extended GHZ states. Clearly, the region shrinks as bipartite entanglement (C_{12}^2) increases. Only small portion of advantage is obtained as C_{12}^2 increases above 0.75 which is shown separately in the figure.

Quantum correlation measure (δ_D) increases monotonically (FIG.6) with that of τ . The degree of violation of 99-th facet inequality (6) varies in a similar fashion like discord monogamy score (δ_D) (for different fixed values of the bipartite entanglement measure C_{12}^2) unlike that of the violation of Svetlichny inequality (2) where the curve shows no such fixed behavior.

IV. RESISTANCE TO NOISE

An interesting observation from both theoretical and physical view point is that of resistance offered to noise by

a pure entangled state. For that, we consider a source \mathcal{S} producing a noisy three qubit state (obtained after pass-




FIG. 6: The degree of violation of 99-th facet inequality (6), Svetlichny inequality (2) and discord monogamy score (δ_D) for subclass S of extended GHZ class are plotted against τ for different fixed values of C_{12}^2 . For τ lying in the range indicated by portion of the τ axis intercepted in between the points where the blue curve and red curve cut the τ axis (other than the origin), nonlocal feature of the corresponding correlations is revealed via violation of 99-th facet inequality (6). For this range of τ , discord monogamy score (δ_D) serves as a better tool for detecting genuine nonlocality compared to Svetlichny inequality (2).

ing the state through a depolarizing channel[72–75]):

$$\omega = \alpha\rho + (1 - \alpha)\frac{\mathbf{I}}{8}, \quad (\text{where } \alpha \in [0, 1]). \quad (24)$$

Here α denotes the visibility of three qubit state ρ which is a measure of the resistance to noise. $1 - \alpha$ denotes the probability with which white noise is introduced in the system by the source. The largest visibility (α) for which ω is local is referred to as the *local visibility threshold* (V). Considering $\rho = |\varphi_{GGHZ}\rangle\langle\varphi_{GGHZ}|$, using 99-th facet inequality one can find $\alpha \in (\frac{3}{1+2\sqrt{1+\tau}}, 1]$ whereas

for Svetlichny ([36]) the range is given by $\alpha \in (\frac{1}{\sqrt{2\tau}}, 1]$. Hence from the point of lowering down the local visibility threshold (V), the efficiency of 99-th facet inequality and Svetlichny (2) depends on the amount of tripartite entanglement present in the corresponding state (produced by the source S) belonging to the GGHZ class (FIG.7). For $\rho = |\chi\rangle\langle\chi|$, a similar analysis (FIG.8) can be made as 99-th facet inequality (6) is violated for $\alpha \in (\frac{3}{1+\sqrt{A+2C}+\sqrt{A-2C}}, 1]$, whereas $\alpha \in (\frac{1}{\sqrt{2\tau+C_{12}^2}}, 1]$, if violation of Svetlichny inequality(2) is considered.




FIG. 7: The difference between the lower bounds α_1 and α_2 of the range of visibility (α) of ω corresponding to the violation of 99-th facet inequality (6) and Svetlichny inequality (2) respectively, is plotted against tripartite entanglement measure (τ) of GGHZ class.

99-th facet inequality gives maximum advantage in terms of lowering down local visibility threshold (V) for the noisy GGHZ state whose corresponding pure counterpart has tripartite entanglement (τ) given by the point A. On the contrary Svetlichny inequality does the same for the noisy state corresponding to the point E.

99-th facet inequality helps in lowering down local visibility (V) of those noisy GGHZ states whose corresponding pure counterparts are characterized by having τ corresponding to any point D lying on the portion of the curve below τ axis.

A. Exposure of individual qubits to noise

Apart from considering a one parameter noisy state, obtained due to the effect of a depolarizing channel on a pure state, it is also interesting to deal with nonlocal character of the states when each of the three subsystems are subjected to:

- independent depolarization with different strength
- individually exposed to amplitude damping with

varying measure of dampness.

1. Independent depolarization

If all the three subsystems of any pure state of the GGHZ class(18) are subjected to depolarization of varying strength then the noisy state is given by:

$$\frac{((J_1 + J_2) \cos^2 \eta + 2J_2 \sin^2 \eta)|000\rangle\langle 000| + ((J_1 + J_2) \sin^2 \eta + 2J_2 \cos^2 \eta)|111\rangle\langle 111| + \frac{(J_1 - J_2)}{2} \sin 2\eta(|000\rangle\langle 111| + |111\rangle\langle 000|)}{J_1 + 3J_2} \quad (25)$$

where $J_1 = (1 - \frac{3p_1}{4})(1 - \frac{3p_2}{4})(1 - \frac{3p_3}{4})$, $J_2 = \frac{1}{64}p_1p_2p_3$ and $p_i (i = 1, 2, 3)$ denoting the strength of the three depolarizing channels through which the 1st, 2nd and 3rd qubit of the state are passed respectively. Here 99th facet in-

equality detects nonlocality of the noisy version of GGHZ state(25) more efficiently than Svetlichny inequality. For instance, with the strength of depolarization of the three channels being $p_1 = 0.8$, $p_2 = 0.7$, $p_3 = 0.6$ and the state




FIG. 8: For any value of τ and C_{12}^2 lying in the shaded region (R), 99-th facet inequality (6) gives advantage over Svetlichny inequality (2), i.e., any pure entangled state (belonging to the subclass S of extended GHZ class), characterized by its tripartite entanglement (τ) and bipartite entanglement content (C_{12}^2) corresponding to a point lying in the region (R), is more resistant to noise when Eq.(6) is considered. On the contrary, for any point lying in the region R' the corresponding pure entangled state offers more resistance to noise via Svetlichny inequality (2) violation. For any value of C_{12}^2 above 0.90 discrete portions of R and R' are obtained (highlighted separately).

parameter being $\eta = 0.69$, 99th facet inequality(6) is violated whereas Svetlichny inequality(2) is not. Similar

conclusion can be drawn for MS subclass(19). For a particular instance for this subclass, the channels used in the previous example can be considered with the state parameter $\eta = 0.69$.

2. Independent amplitude damping

For practical purposes, analysis of dissipation of energy from a quantum system is an important application of quantum operations. In that context amplitude damping plays a significant role in characterizing the general behavior of some related physical processes. For instance, it may be used to describe the dynamics of an atom which is spontaneously releasing photon, dynamics of an excited spin system tending towards equilibrium with its environment or analyzing state of a photon when subjected to scattering and attenuation in a cavity or interferometer. Mathematically, amplitude damping is represented using Krauss operator formalism [72, 76]. When a two qubit state ρ is passed through an amplitude damping channel, the resulting noisy state is given by

$$\theta_{AD}(\rho) = E_0 \rho E_0^\dagger + E_1 \rho E_1^\dagger \quad (26)$$

where $E_0 = \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{1-\gamma} \end{pmatrix}$ and $E_1 = \begin{pmatrix} 0 & \sqrt{\gamma} \\ 0 & 0 \end{pmatrix}$ are the Krauss operators used for representing the amplitude damping operation. All the three qubits of the state are passed in order through three different amplitude damping channel having varying capacity characterized by the parameters γ_i ($i = 1, 2, 3$), precisely i th qubit passes through channel characterized by γ_i ($i = 1, 2, 3$). In such a situation, noisy version of any state of the GHZ class is given by:

$$\frac{\cos^2 \eta |000\rangle\langle 000| + \left(\frac{D_1 + D_2}{2}\right) |000\rangle\langle 111| + \left(\frac{D_1 \sin 2\eta}{2}\right) |111\rangle\langle 000| + D_1^2 \sin^2 \eta |111\rangle\langle 111|}{\cos^2 \eta + D_1^2 \sin^2 \eta} \quad (27)$$

with $D_1 = \sqrt{(1-\gamma_1)(1-\gamma_2)(1-\gamma_3)}$ and $D_2 = \gamma_1 \gamma_2 \gamma_3$. As in the case of depolarizing channel, in this case also 99th facet inequality emerges as a better inequality compared to svetlichny(2) to detect nonlocal behavior of any state belonging to this class. For a particular instance, one can consider 1st, 2nd and 3rd qubit of the state $0.995|000\rangle + 0.099|111\rangle$ passing through channels with $\gamma_1 = 0.1$, $\gamma_2 = 0.08$, $\gamma_3 = 0.09$ respectively. For MS subclass also it can be concluded that 99th facet inequality(6) surpasses Svetlichny inequality(2) due to its efficiency to detect genuine nonlocality of the noisy states belonging to this subclass. For a particular numerical example, one may consider the state $\frac{1}{\sqrt{2}}(|000\rangle + 0.955|110\rangle + 0.296|111\rangle)$ and three amplitude damping channels with $\gamma_1 = 0.33$, $\gamma_2 = 0.15$, $\gamma_3 = 0.09$.

V. EXPERIMENTAL DETECTION OF GENUINE NONLOCALITY

In recent times high fidelity tripartite genuine entangled states are generated experimentally([51],[52],[53],[54]). For instance, in [52], an experimental set up is reported where maximally entangled GHZ state is generated with 86.2% fidelity. So in any such experiment generating genuine tripartite entangled states, the nonlocal behavior of the generated state can be detected by using suitable tripartite Bell inequality. As already discussed before, Svetlichny nonlocality lacks proper physical interpretation, to be more precise there exists no clear indication whether signaling is allowed or not while constructing S_2 correlations

and even if its existence is assumed, then the direction of signaling(one way or both way) cannot be properly interpreted. Due to this sort of ambiguity in definition of S_2 correlations, Svetlichny inequality becomes unfit to be used to detect genuine nonlocality of any experimentally generated genuine entangled state. On the contrary, while framing 99th facet inequality no signaling is assumed. Hence definition of NS_2 correlations free from this sort of ambiguity. This in turn points out that the correlation statistics generated while constructing the genuine entangled state experimentally may be used to test the nonlocal nature of the state via the 99th facet inequality. Apart from the physical point of view, theoretically we have shown that 99th facet inequality can detect genuine nonlocality of any state belonging to the GHZ class unlike that of Svetlichny inequality. This in turn gives advantage for experimental purposes. To be more precise, if the state $0.966|000\rangle + 0.259|111\rangle$ is generated experimentally, then Svetlichny inequality fails to detect genuine nonlocality(16) of this genuine tripartite entangled state($\tau = 0.25 < \frac{1}{3}$) whereas 99th facet can detect its nonlocal behavior from the correlation statistics(15).

VI. FAMILY OF SOME HIGHER RANK MIXED STATES

For a three-qubit mixed state, $\rho = \sum_i p_i \rho_i$ analytical expressions of three-tangle are given in terms of convex roof: $\tau(\rho) = \min_{p_i} \sum_i p_i \tau(\rho_i)$. So far the analytical expressions for the three-tangle (τ) for all states of rank-2 [55, 56], rank-3, 4 [57, 58] and rank- n where $n = 5, 6, 7, 8$ [59] are reported. Mixture of GHZ and W states and that of GHZ, W and flipped W states are mixed states of rank-2 and rank-3 respectively:

$$\rho_2 = p|GHZ\rangle\langle GHZ| + (1-p)|W\rangle\langle W| \quad (28)$$

$$\rho_3^k = p|GHZ\rangle\langle GHZ| + q|W\rangle\langle W| + (1-p-q)|\widetilde{W}\rangle\langle\widetilde{W}| \quad (29)$$

where $|GHZ\rangle = \frac{|000\rangle + |111\rangle}{\sqrt{2}}$, $|W\rangle = \frac{|001\rangle + |010\rangle + |100\rangle}{\sqrt{3}}$, $|\widetilde{W}\rangle = \frac{|011\rangle + |110\rangle + |101\rangle}{\sqrt{3}}$. and $q = \frac{1-p}{k}$, k being any positive integer. For $k = 1$, $\rho_3^k = \rho_2$. The study of three-tangle (τ) of these mixed states focusses on the characterization of genuine tripartite entanglement. But none of these studies demonstrates the nonlocal behavior of these states. The range of p for which rank-2 and rank-3 (for $k = 2, 3, 10$) mixed states exhibit nonlocality via violation of 99-th facet (6) and Svetlichny inequality(2) is summarised in Table I.

As pointed out in [59], the families of high rank ($n = 5, \dots, 8$) mixed states are interesting for practical purposes as their tripartite entanglement features would reveal further relations between the quantum phase transitions and quantum entanglement just like concurrence of mixed two-qubit states which have been applied in the study of quantum phase transitions. Besides, as τ , being

State	$\tau > 0$	Violation of Eq.(6)	Violation of Eq.(2)
ρ_2	$p \geq 0.6268$	$p \geq 0.811876$	$p \geq 0.707109$
ρ_3^2	$p \geq 0.75$	$p \geq 0.819964$	$p \geq 0.70719$
ρ_3^3	$p \geq 0.7452$	$p \geq 0.818825$	$p \geq 0.707109$
ρ_3^{10}	$p \geq 0.7452$	$p \geq 0.814789$	$p \geq 0.707109$

TABLE I: The table gives a comparison between the bounds for violation of Svetlichny inequality and 99th facet inequality along with that of the measure of tripartite entanglement(τ) for some of the mixed states ρ_2, ρ_3^i ($i=2,3,10$).

a measure of genuine tripartite entanglement, is capable of giving certain information in the scheme of quantum copy machine or three-party quantum teleportation [77]. So a systematic study dealing with the connection of this measure of genuine entanglement with that of violation of a Bell-type inequality (facet of a local polytope) can be useful in related research works. The families of rank n ($4, \dots, 8$) are [58, 59]:

$$\rho_4 = p|\Lambda, 1+\rangle\langle\Lambda, 1+| + \frac{(1-p)}{3}\Pi \quad (n=4) \quad (30)$$

$$\rho_5 = p|\Lambda, 1+\rangle\langle\Lambda, 1+| + \frac{(1-p)}{10}(|\Lambda, 1-\rangle\langle\Lambda, 1-| + 3\Pi) \quad (n=5) \quad (31)$$

$$\rho_6 = p|\Lambda, 2-\rangle\langle\Lambda, 2-| + \frac{(1-p)}{11}(\Omega + 3\Pi) \quad (n=6) \quad (32)$$

$$\rho_7 = p|\Lambda, 3-\rangle\langle\Lambda, 3-| + \frac{(1-p)}{34}(|\Lambda, 2-\rangle\langle\Lambda, 2-| + 3\Omega + 9\Pi) \quad (n=7) \quad (33)$$

$$\rho_8 = p|\Lambda, 4-\rangle\langle\Lambda, 4-| + \frac{(1-p)}{35}(|\Lambda, 2-\rangle\langle\Lambda, 2-| + |\Lambda, 3-\rangle\langle\Lambda, 3-| + 3\Omega + 9\Pi) \quad (n=8) \quad (34)$$

where $|\Lambda, 1\pm\rangle = \frac{|000\rangle \pm |111\rangle}{\sqrt{2}}$, $|\Lambda, 2\pm\rangle = \frac{|110\rangle \pm |001\rangle}{\sqrt{2}}$, $|\Lambda, 3\pm\rangle = \frac{|101\rangle \pm |010\rangle}{\sqrt{2}}$, $|\Lambda, 4\pm\rangle = \frac{|011\rangle \pm |100\rangle}{\sqrt{2}}$, $\Omega = |\Lambda, 1+\rangle\langle\Lambda, 1+| + |\Lambda, 1-\rangle\langle\Lambda, 1-|$ and $\Pi = |\Lambda, 2+\rangle\langle\Lambda, 2+| + |\Lambda, 3+\rangle\langle\Lambda, 3+| + |\Lambda, 4+\rangle\langle\Lambda, 4+|$.

The closed form of the bounds (B_6 and B_7) of 99-th facet

inequality for ρ_4 and ρ_5 are respectively:

$$B_6 = \frac{2\sqrt{16p^2 - 8p + 10} + |1 - 4p|}{3} \quad (35)$$

$$B_7 = \frac{2\sqrt{37p^2 - 4p + 17} + |1 - 6p|}{5}. \quad (36)$$

ρ_4 violates Eq.(6) for $p \geq 0.726$ and Svetlichny inequality (2) for $p \geq 0.72$, whereas it has a nonzero tangle for $p \geq 0.75$ [58]. As genuine multipartite nonlocal nature of the correlations produced by a state is a signature of existence of genuine multipartite entanglement of the state, the 185-th facet (i.e., Svetlichny inequality (2)) can be regarded as a better measure of genuine tripartite entanglement of ρ_5 than the three tangle(τ). A similar sort of results holds for ρ_5 which violates Eq.(2) and Eq.(6) for $p \geq 0.710858$ and $p \geq 0.729157$ respectively, but has $\tau > 0$ for $p \geq 0.737$ [59]. Hence for $p \in [0.710858, 0.737]$ violation of a facet inequality of

NS_2 local polytope serves as a better measure for detecting genuine tripartite entanglement compared to three tangle. However, a comparative study of the range of p for non zero three tangle and that of 99-th (6) and 185-th (2) facet violation for the other three families of higher rank mixed states ($n = 6, 7, 8$) does not yield analogous result and so these facet inequalities can only be used to demonstrate the genuine tripartite nonlocality of the correlations produced in a system using these families of mixed states upto projective measurements. The results obtained are summarized as follows(Table II):

State	Bound for Eq.(6)	$\tau > 0$	Violation of Eq.(6)	Violation of Eq.(2)
ρ_6	$\frac{\sqrt{(1+10p)^2+(6(1-p)+ 3-14p)^2+(12p-1)}}{11}$	$p \geq 0.2143$	$p \geq 0.756458$	$p \geq 0.765134$
ρ_7	$\sqrt{(-.11765 + 1.11765p)^2 + B^2} + \sqrt{(.11765 + 0.8824p)^2 + B^2} + 0.0588 + 0.9412p$, $B = \frac{1-p}{2} + 0.26470 - 1.26470p $	$p \geq 0.2062$	$p \geq 0.759185$	$p \geq 0.76444$
ρ_8	$\sqrt{(0.0857 + 0.9143p)^2 + C^2} + \sqrt{(-0.1428 + 1.1429p)^2 + C^2} + 0.0857 + 0.9142p$, $C = 0.4572(1-p) + .2571 - 1.2571p $	$p \geq 0.2490$	$p \geq 0.75843$	$p \geq 0.763645$

TABLE II: The table gives a comparison between the bounds for violation of Svetlichny inequality and 99th facet inequality along with that of the measure of tripartite entanglement(τ) for some of the higher rank mixed states ρ_6 , ρ_7 and ρ_8 .

VII. CONCLUSION:

In conclusion, the above systematic study exploiting the relation between genuine tripartite entanglement measure and that of genuine tripartite nonlocality reveals the efficiency of a facet (99-th) of NS_2 local polytope as a tripartite Bell-type inequality over Svetlichny inequality (which is reported as the 185-th facet of NS_2 local polytope) for detecting genuine three way nonlocality of GGHZ and a subclass S of extended GHZ class of pure tripartite genuine entangled states. This in turn helps to prove the conjecture made by Bancal in([31]) for the GGHZ class of states. Apart from the obvious changes in the analytical expressions of the bounds of 99-th facet inequality from that of Svetlichny inequality (2) for the corresponding class of states, it is interesting to note that the rectification (made by Bancal in ([31])), of local-nonlocal hybrid correlations (1) and thereby giving a more physically well motivated definition of the same (1,4,5), contributes in developing more efficient measures for demonstrating three way nonlocality. So far this sort of analysis has been reported only in continuous variable systems of Gaussian states ([44]). A comparative study between discord monogamy score and violation of these two facet inequalities clearly points out that from the perspective of revealing quantumness in a multipartite

system, 99-th facet inequality (6) surpasses Svetlichny inequality (2). Besides, in the context of analyzing the interplay between mixed state genuine entanglement and the nonlocality of the corresponding correlations emerging in a quantum system, Eq.(6) gives advantage over Svetlichny inequality (2) in many cases such as lowering down the local visibility threshold V of a subclass of noisy GGHZ and subclass S of extended GHZ states and also for some family of higher rank mixed states. Here it is worth mentioning that for some family of mixed states of rank-4 and rank-5, these facet inequalities (both 2 and 6) can even be used as a better measure of genuine tripartite entanglement over the usual measure of three tangle (τ). It will be interesting to explore further in this approach that whether one can develop a more efficient measure of detecting three way nonlocality (both mixed and pure) and also if possible a better measure of genuine entanglement for other family of mixed states by choosing a suitable facet of the NS_2 local polytope and also by performing more general measurements which in turn may lead to prove the conjecture made by Bancal et.al. ([31]). Apart from this, there are many other future directions. The analysis made here can be used in related physical experiments. One may also explore the relation between tripartite information content of GGHZ class [78] and the corresponding nonlocal correlations. There exists close correspondence between Bell inequal-

ity violation and nonlocal games [79]. So from that perspective it will be interesting to develop a detailed study focussing on the efficiency of GGHZ and subclass S of extended GHZ class in various protocols based on nonlocal games [80].

Acknowledgement: The authors acknowledge fruitful

discussions with Mr. A.Sen and Ms. S.Karmakar. The author KM acknowledges financial support from University Grants Commission(UGC), New Delhi and the author D. Sarkar acknowledges the support from DST, India.

-
- [1] A. Acín, A. Andrianov, L. Costa, E. Jané, J. I. Latorre and R. Tarrach, *Phys. Rev. Lett.* **85**, 7 (2000).
- [2] J. S. Bell, *Physics* (Long Island City, N.Y.) **1**, 195(1964).
- [3] J. S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, England,1988).
- [4] M. Genovese, *Physics Reports* **413**, 319 (2005).
- [5] A. Aspect, *Nature* **398**, 189 (1999).
- [6] M. Seevinck and J. Uffink *Phys. Rev. A* **78**, 032101 (2008).
- [7] M. Hein, J. Eisert and H. J. Briegel, *Phys. Rev. A* **69**, 062311 (2004).
- [8] K. Chen and Lo H-K, *Quantum Inf. Comput.* **7**, 689–715 (2007).
- [9] V. Scarani and N. Gisin, *Phys. Rev. Lett.* **87**, 117901 (2001).
- [10] M. Seevinck, Parts and wholes, An inquiry into quantum and classical correlations, arXiv:0811.1027 (2008).
- [11] P. Zoller et al, *Eur. Phys. J. D* **36**, 203–28 (2005).
- [12] C-Y Lu , X-Q Zhou, O. Guhne, W-B Gao, J Zhang, Z-S Yuan, A Goebel, T Yang and J-W Pan, *Nature Phys.* **3**, 91–5 (2007).
- [13] R. Laflamme, E. Knill, W. Zurek, P. Catasti and S. V. S. Mariappan, *Phil. Trans. R. Soc. A* **356**, 1941–8 (1998).
- [14] D. Bouwmeester, J-W. Pan, V. Daniell, H. Weinfurter and A. Zeilinger, *Phys. Rev. Lett.* **82**, 1345–9 (1999).
- [15] C. F. Roos, M. Riebe, H. Haffner, W. Hansel, J. Benhelm, G. P. T. Lancaster, C. Becher, F. Schmidt-Kaler and R. Blatt, *Science* **304**, 1478–80 (2004).
- [16] R. Cleve and H. Buhrman, *Phys. Rev. A* **56**, 1201 (1997).
- [17] J. Barrett, L. Hardy, and A. Kent, *Phys. Rev. Lett.* **95**, 010503 (2005).
- [18] Ll. Masanes, *Phys. Rev. Lett.* **102**, 140501 (2009).
- [19] A. Acin, N. Brunner, N. Gisin, S. Massar, S. Pironio, and V. Scarani, *Phys. Rev. Lett.* **98**, 230501 (2007);
- [20] Ll. Masanes, S. Pironio, A. Acin, *Nature Comm.* **2**, 238 (2011).
- [21] S. Pironio et al, *Nature* **464**, 1021 (2010).
- [22] R. Colbeck and A. Kent, *J. Phys. A: Math. and Theor.* **44**, 095305 (2011).
- [23] R. Raussendorf and H. J. Briegel, *Phys. Rev. Lett.* **86**, 5188 (2001).
- [24] R. Raussendorf, D. E. Browne and H. J. Briegel, *Phys. Rev. A* **68**, 022312 (2003).
- [25] T. Roscilde, P. Verrucchi, A. Fubini, S. Haas and V. Tognetti, *Phys. Rev. Lett.* **94**, 147208 (2005).
- [26] C. Brukner, V. Vedral and A. Zeilinger, *phys. Rev. A* **73**, 012110 (2006).
- [27] A. Osterloh, L. Amico, G. Falci and R. Fazio, *Nature* **416**, 608–10 (2002).
- [28] G. Vidal, J. I. Latorre, E. Rico and A. Kitaev, *Phys. Rev. Lett.* **90**, 227902 (2003).
- [29] T. J. Osborne and M. A. Nielsen, *Phys. Rev. A* **66**, 032110 (2002).
- [30] G. Svetlichny, *Phys. Rev. D* **35**, 3066 (1987).
- [31] J.-D. Bancal, J. Barrett, N. Gisin, and S. Pironio, *Phys. Rev. A* **88**, 014102 (2013).
- [32] D. Collins, N. Gisin, S. Popescu, D. Roberts, and V. Scarani, *Phys. Rev. Lett.* **88**, 170405 (2002).
- [33] M. Seevinck and G. Svetlichny, *Phys. Rev. Lett.* **89**, 060401 (2002).
- [34] J.-D. Bancal, N. Brunner, N. Gisin, and Y.-C. Liang, *Phys. Rev. Lett.* **106**, 020405 (2011).
- [35] J.-L. Chen, D.-L. Deng, H.-Y. Su, C. Wu, and C.H. Oh, *Phys. Rev. A* **83**, 022316 (2011).
- [36] S. Ghose, N. Sinclair, S. Debnath, P. Rungta, and R. Stock, *Phys. Rev. Lett.* **102**, 250404 (2009).
- [37] A. Ajoy and P. Rungta, *Phys. Rev. A* **81**, 052334 (2010).
- [38] J. Lavoie, R. Kaltenbaek, and K. J. Resch, *New J. Phys.* **11**, 073051 (2009).
- [39] R. Gallego, L. E. Würflinger, A. Acin, and M. Navascues, *Phys. Rev. Lett.* **109**, 070401 (2012).
- [40] R. Gallego, L. E. Würflinger, A. Acin, and M. Navascues, *Phys. Rev. Lett.* **90**, 227902 (2003).
- [41] V. Coffman, J. Kundu, and W. K. Wootters, *Phys. Rev. A* **61**, 052306 (2000).
- [42] W. Dur, G. Vidal and J. I. Cirac, *Phys. Rev. A* **62** 062314 (2000).
- [43] H. A. Carteret and A. Sudbery, *J. Phys. A: Math. Gen.* **33**, 4981–5002 (2000).
- [44] G. Adesso and S. Piano, *Phys. Rev. Lett.* **112**, 010401 (2009).
- [45] ML. Almeida, D. Cavalcanti, V. Scarani and A. Acin, *Phys. Rev. A* **81**, 052111 (2010).
- [46] R. Rahaman, M. Wiesniak, and M. Zukowski, arXiv:1303.0128.
- [47] Q. Chen, S. Yu, C. Zhang, C.H. Lai, and C.H. Oh, *Phys. Rev. Lett.* **112**, 140404 (2012).
- [48] S. Yu, and C.H. Oh, arXiv:1306.5330.
- [49] R. Prabhu, A.K. Pati, A. Sen(De), and U. Sen, *Phys. Rev. A* **85**, 040102 (R) (2012).
- [50] R. Prabhu, A.K. Pati, A. Sen(De), and U. Sen, *Phys. Rev. A* **86**, 052337 (2012).
- [51] L. DiCarlo, M. D. Reed, L. Sun, B. R. Johnson, J.M. Chow, J. M. Gambetta, L. Frunzio, S. M. Girvin, M. H. Devoret and R. J. Schoelkopf, *Nature*, **09416**, 574-578, (2010).
- [52] D. R. Hamel, L. K. S. H. Hubel, A. J. Miller, F. Marsili, V. B. Verma, R. P. Mirin, S. W. Nam, K. J. Resch and T. Jennewein, arXiv:1404.7131.
- [53] J. Lavoie, R. Kaltenbaek, K. J. Resch, *New Journal of Physics*, **11**, 073051 (2009).
- [54] H.-X. Lu, et al., *Phys. Rev. A* **84**, 012111 (2011).
- [55] R. Lohmayer, A. Osterloh, J. Siewert, and A. Uhlmann, *Phys. Rev. Lett.* **97**, 260502 (2006).
- [56] C. Eltschka, A. Osterloh, J. Siewert, and A. Uhlmann,

New J.Phys. **10**, 043014 (2008).

- [57] E. Jung, M.R. Hwang, D.K. Park, and J.W. Son, Phys. Rev. A **79**, 024306 (2009).
- [58] E. Jung, D. Park, and J.W. Son, Phys. Rev. A **80**, 010301 (2009).
- [59] S-J. He, X-H. Wang, S-M. Fei, H-X. Sun and Q-Y. Wen, Commun. Theor. Phys. **55**, 251-256 (2011).
- [60] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
- [61] L. Henderson and V. Vedral, J. Phys. A **34**, 6899 (2001).
- [62] H. Ollivier and W.H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001).
- [63] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, Rev. Mod. Phys. **84**, 1655-1707 (2012).
- [64] D. M. Greenberger, M. Horne, A. Shimony, and A. Zeilinger, Am. J. Phys. **58**, 1131 (1990).
- [65] N. Linden, S. Popescu and W. K. Wootters, Phys. Rev. Lett. **89**, 207901 (2002).
- [66] R. J. Nelson, D. G. Cory, S. Lloyd, Phys. Rev. A **61**, 022106 (2000).
- [67] C. Emary and C. W. J. Beenakker, Phys. Rev. A **69**, 032317 (2004).
- [68] S. Ghose, S. Debnath, N. Sinclair, A. Kabra and R. Stock, J. Phys. A: Math. Theor. **43**, 445301(2010).
- [69] J. F. Clauser, et al., Phys. Rev. Lett. **23**, 880 (1969).
- [70] N. Gisin, Phys. Lett. A **154**, 201 (1991).
- [71] S. Popescu and D. Rohrlich, Phys. Lett. A **166**, 293 (1992).
- [72] M. A. Nielsen and I. L. Chuang, "Quantum computation and quantum information", Cambridge University Press (2000).
- [73] C.H Bennet, D.P. DiVincenzo, J. A. Smolin, Phys. Rev. Lett. **78**, 3217 (1997).
- [74] C.H Bennet, P. W. Shor, J. A. Smolin and A. V. Thapliyal, Phys. Rev. Lett. **83**, 3081 (1999).
- [75] D. Bruss, L. Faoro, C. Macchiavello, and G. M. Palma, Journal of Modern Optics, **47**, 325 (2000).
- [76] K. Kraus, States, Effects and operations (Springer-Verlag, Berlin, 1983).
- [77] A. Karlsson and M. Bourennane, Phys. Rev. A **58**, 4394 (1998).
- [78] N. Linden, S. Popescu and W. K. Wootters, Phys. Rev. Lett. **89**, 207901 (2002).
- [79] J. Silman, S. Machnes and N. Aharon, Phys. Lett. A **372**, 3796-800 (2008).
- [80] R. Cleve, P. Høyer, B. Toner, and J. Watrous, Proceedings of the 19th IEEE Conference on Computational Complexity (2004), p. 236.
- [81] W.H. Zurek, in Quantum Optics, Experimental Gravitation and Measurement Theory, eds. P. Meystre and M.O. Scully (Plenum, New York, 1983)
- [82] S.M. Barnett and S.J.D. Phoenix, Phys. Rev. A **40**, 2404 (1989).
- [83] N.J. Cerf and C. Adami, Phys. Rev. Lett. **79**, 5194 (1997).
- [84] B. Schumacher and M.A. Nielsen, Phys. Rev. A **54**, 2629 (1996).
- [85] B. Groisman, S. Popescu, and A. Winter, *ibid.* **72**, 032317 (2005).
- [86] Q. Chen, C. Zhang, S. Yu, X. X. Yi, and C. H. Oh, Phys. Rev. A **84**, 042313 (2011).

VIII. APPENDIX A

In order to obtain the bound given in (14), we consider the following measurements: $X = \vec{x} \cdot \vec{\sigma}_1$ or, $\hat{X} = \vec{x} \cdot \vec{\sigma}_1$ on qubit 1, $Y = \vec{y} \cdot \vec{\sigma}_2$ or, $\hat{Y} = \vec{y} \cdot \vec{\sigma}_2$ on qubit 2, and $Z = \vec{z} \cdot \vec{\sigma}_3$ or, $\hat{Z} = \vec{z} \cdot \vec{\sigma}_3$ on qubit 3, where $\vec{x}, \vec{y}, \vec{z}$ and \vec{x}, \vec{z} are unit vectors and σ_i are the spin projection operators that can be written in terms of the Pauli matrices. Representing the unit vectors in spherical coordinate, we have, $\vec{x} = (\sin \theta_{a_0} \cos \phi_{a_0}, \sin \theta_{a_0} \sin \phi_{a_0}, \cos \theta_{a_0})$, $\vec{y} = (\sin \alpha b_0 \cos \beta b_0, \sin \alpha b_0 \sin \beta b_0, \cos \alpha b_0)$ and $\vec{z} = (\sin \zeta c_0 \cos \eta c_0, \sin \zeta c_0 \sin \eta c_0, \cos \zeta c_0)$ and similarly, we define, \vec{x}, \vec{y} and \vec{z} by replacing 0 in the indices by 1. Then the expectation value of the operator NS (6) with respect to the state $|\varphi_{GGHZ}\rangle$ gives

$$NS(|\varphi_{GGHZ}\rangle) = \cos(\theta_{a_0}) \cos(\alpha b_0) (\cos(\zeta c_0) \cos(2\eta) - \cos(\zeta c_1) \cos(2\eta)) + \sin(\theta_{a_0}) \sin(\alpha b_0) (\sin(\zeta c_0) \cos(\beta b_0 + \eta c_0 + \phi_{a_0}) \sin(2\eta) - \sin(\zeta c_1) \cos(\beta b_0 + \eta c_1 + \phi_{a_0}) \sin(2\eta)) + \cos(\zeta c_0) \cos(\alpha b_1) + \cos(\alpha b_1) \cos(\theta_{a_1}) + \cos(\zeta c_1) \cos(\theta_{a_1}). \quad (37)$$

Hence in order to get a maximum value of the NS operator defined in (6), we have to perform a maximization over 12 measurement angles. We first find the global maximum of $NS(|\varphi_{GGHZ}\rangle)$ with respect to θ_{a_0} and αb_0 . We begin with by finding all critical points of $NS(|\varphi_{GGHZ}\rangle)$ inside the region $R = [0, 2\pi] \times [0, 2\pi]$ which are namely $(0, 0)$, $(\frac{\pi}{2}, -\frac{\pi}{2})$, $(-\frac{\pi}{2}, \frac{\pi}{2})$ and $(\frac{\pi}{2}, \frac{\pi}{2})$. Among all these critical points the point $(\frac{\pi}{2}, \frac{\pi}{2})$ gives the global maximum of $NS(|\varphi_{GGHZ}\rangle)$. Thus, we have,

$$NS(|\varphi_{GGHZ}\rangle) \leq \sin(\zeta c_0) \cos(\beta b_0 + \eta c_0) + \phi_{a_0} \sin(2\eta) - \sin(\zeta c_1) \cos(\beta b_0 + \eta c_1 + \phi_{a_0}) \sin(2\eta) + \cos(\zeta c_0) \cos(\alpha b_1) + \cos(\zeta c_1) \cos(\theta_{a_1}) + \cos(\alpha b_1) \cos(\theta_{a_1}) \quad (38)$$

which when maximized with respect to ζc_0 and ζc_1 , gives

$$NS(|\varphi_{GGHZ}\rangle) \leq \sqrt{\cos^2(\alpha b_1) + \sin^2(2\eta)} + \sqrt{\cos^2(\theta_{a_1}) + \sin^2(2\eta)} + \cos(\alpha b_1) \cos(\theta_{a_1}). \quad (39)$$

The last inequality is obtained by using the inequality $x \cos \theta + y \sin \theta \leq \sqrt{x^2 + y^2}$ (the equality holds for $\tan \theta = \frac{y}{x}$) and by letting $\cos^2(\beta b_0 + \eta c_0 + \phi_{a_0}) = 1$, $\cos^2(\beta b_0 + \eta c_1 + \phi_{a_0}) = 1$. Again by classifying all the critical points, it is easy to check that the expression in Eq.(39) gives the global maximum with respect to αb_1 and θ_{a_1} for the critical point $(0, 0)$. Therefore, Eq.(39) reduces to:

$$NS(|\varphi_{GGHZ}\rangle) \leq 1 + 2\sqrt{1 + \sin^2 2\eta} \quad (40)$$

as stated in Eq.(14).

IX. APPENDIX B

For extended GHZ states, the expression of NS (6) is:

$$NS(|\chi\rangle) = A_0((1 - 2\lambda_4^2)Y_1 - 2\lambda_3\lambda_4 \sin(\zeta c1) \cos(\eta c1)) + D_0(2\lambda_0\lambda_3Y_1B_0 + 2\lambda_0\lambda_4 \sin(\zeta c0)C_0 - 2\lambda_0\lambda_4 \sin(\zeta c1)C_1) \\ + Y_0(1 - 2\lambda_3^2) - 2\lambda_3\lambda_4 \sin(\zeta c0) \cos(\eta c0) \cos(\alpha b1) - 2\lambda_3\lambda_4 \sin(\zeta c1) \cos(\eta c1) \cos(\theta a1) + A_1 + 2\lambda_0\lambda_3B_1D_1. \quad (41)$$

where $X_i = \cos(\zeta ci)$, $A_i = \cos(\alpha bi) \cos(\theta ai)$, $B_i = \cos(\beta bi + \phi ai)$, $C_i = \cos(\beta b0 + \phi a0 + \eta ci)$, $D_i = \sin(\alpha b0) \sin(\theta a0)$ ($i = 0, 1$) and $X_0 + X_1 = Y_0$, $X_0 - X_1 = Y_1$

Considering above expression as a function of $\sin(\theta a0)$ and $\sin(\alpha b0)$, we get the critical points: $(0, 0)$, $(1, 1)$,

$(1, -1)$, $(-1, 1)$, $(-1, -1)$. The global maximum of this function gives the required bound.

Case1: For $\sin(\theta a0) = 0$ and $\sin(\alpha b0) = 0$

$$NS(|\chi\rangle) = X_0(T_1 \cos(\alpha b1) + T_2) + \sin(\zeta c0)(-2T_3 \cos(\eta c0) \cos(\alpha b1) + 2T_3 \cos(\eta c0))X_1(T_1 \cos(\theta a1) \\ - T_2) + \sin(\zeta c1)(-2T_3 \cos(\eta c1) \cos(\theta a1) - 2T_3 \cos(\eta c1)) + A_1 + 2\lambda_0\lambda_3B_1D_1 \\ \leq \sqrt{\cos^2(\alpha b1)(T_1^2 + (2T_3 \cos(\eta c0))^2) + 2(T_1T_2 - 4(T_3)^2) \cos(\alpha b1) + (2T_3 \cos(\eta c0))^2} + \\ \sqrt{\cos^2(\theta a1)(T_1^2 + (2T_3 \cos(\eta c1))^2) + 2(T_1T_2 - 4(T_3)^2) \cos(\theta a1) + (2T_3 \cos(\eta c1))^2} + A_1 + 2\lambda_0\lambda_3B_1D_1.$$

where $T_1=(1 - 2\lambda_3^2)$, $T_2=(1 - 2\lambda_4^2)$ and $T_3=\lambda_3\lambda_4$ The above expression can be considered as a function of $\cos(\alpha b1)$ and $\cos(\theta a1)$ which is maximum for $\cos(\alpha b1) = 0$ and $\cos(\theta a1) = 0$. Hence the bound gets modified as

$$\sqrt{T_1^2 + T_2^2 + 2T_1T_2} + \sqrt{T_1^2 + T_2^2 - 2T_1T_2 + (4T_3 \cos(\eta c1))^2} + 1.$$

which is maximum for $\eta c1 = 0$. So, we get the bound as,

$$\sqrt{4\lambda_0^4} + \sqrt{4(\lambda_3^2 - \lambda_4^2)^2 + 16\lambda_3^2\lambda_4^2} + 1$$

which on further simplification gives the value 3.

Case2: For $\sin(\theta a0) = 1$ and $\sin(\alpha b0) = -1$

$$NS(|\chi\rangle) = X_0(-2\lambda_0\lambda_3B_0 + T_1 \cos(\alpha b1)) + \sin(\zeta c0)(-2\lambda_0\lambda_4C_0 - 2T_3 \cos(\eta c0) \cos(\alpha b1)) + \cos(\zeta c1)(-2\lambda_0\lambda_3B_0 + T_1 \cos(\theta a1)) \\ + \sin(\zeta c1)(-2\lambda_0\lambda_4C_1 - 2T_3 \cos(\eta c1) \cos(\theta a1)) + A_1 + 2\lambda_0\lambda_3B_1D_1 \\ \leq \sqrt{(2\lambda_0\lambda_3B_0 - T_1 \cos(\alpha b1))^2 + (-2\lambda_0\lambda_4C_0 - 2T_3 \cos(\eta c0) \cos(\alpha b1))^2} \\ + \sqrt{(2\lambda_0\lambda_3B_0 + T_1 \cos(\theta a1))^2 + (2\lambda_0\lambda_4C_1 - 2T_3 \cos(\eta c1) \cos(\theta a1))^2} \\ = G(\cos(\alpha b1), \cos(\theta a1))$$

which is maximum for $\cos(\alpha b1) = 0$ and $\cos(\theta a1) = 0$. Thus, the bound modified as:

$$\sqrt{(2\lambda_0\lambda_3B_0 - T_1)^2 + (2\lambda_0\lambda_4C_0 + 2T_3 \cos(\eta c0))^2}$$

$$+ \sqrt{(2\lambda_0\lambda_3B_0 + T_1)^2 + (2\lambda_0\lambda_4C_1 - 2T_3 \cos(\eta c1))^2}.$$

Putting $\beta b0 = \phi a0 = 0$, and then maximizing over $\eta c0$ and $\eta c1$ (maximum for $\eta c0 = \eta c1 = 0$), we get the bound:

$$\sqrt{(2\lambda_0\lambda_3 - T_1)^2 + (2\lambda_0\lambda_4 + 2T_3)^2} + \sqrt{(2\lambda_0\lambda_3 + T_1)^2 + (2\lambda_0\lambda_4 - 2T_3)^2}$$

This on further simplification gives

$$1 + \sqrt{1 + 4\lambda_0^2\lambda_4^2 + 4\lambda_0\lambda_3(1 - 2\lambda_0^2)} \\ + \sqrt{1 + 4\lambda_0^2\lambda_4^2 - 4\lambda_0\lambda_3(1 - 2\lambda_0^2)}.$$

So, $f(1, -1) = 1 + \sqrt{1 + 4\lambda_0^2\lambda_4^2 + 4\lambda_0\lambda_3(1 - 2\lambda_0^2)} + \sqrt{1 + 4\lambda_0^2\lambda_4^2 - 4\lambda_0\lambda_3(1 - 2\lambda_0^2)}$.

Similarly, it can be shown that $f(-1, 1) = f(1, 1) = f(-1, -1) = f(1, -1)$.

Using $\tau = 4\lambda_0^2\lambda_4^2$ and $C_{12}^2 = 4\lambda_0^2\lambda_3^2$ we get: $(1 - 2\lambda_0^2)^2 = 1 - (\tau + C_{12}^2)$.

Hence

$$f(1, -1) = 1 + \sqrt{1 + \tau + 2\sqrt{C_{12}^2(1 - (\tau + C_{12}^2))}} \\ + \sqrt{1 + \tau - 2\sqrt{C_{12}^2(1 - (\tau + C_{12}^2))}}.$$

Clearly,

$$f(1, -1) > 3 \text{ if } \sqrt{1 + \tau + 2\sqrt{C_{12}^2(1 - (\tau + C_{12}^2))}} \\ + \sqrt{1 + \tau - 2\sqrt{C_{12}^2(1 - (\tau + C_{12}^2))}} > 2$$

which on simplification gives, $f(1, -1) > 3$ if $\tau > C_{12}^2 - \tau C_{12}^2 - C_{12}^4$, i.e., $\tau > \frac{C_{12}^2(1 - C_{12}^2)}{1 + C_{12}^2}$. So, the global maximum of $f(\sin(\theta a0), \sin(\theta b0))$ is given by

$$\max\{3, 1 + \sqrt{1 + \tau + 2\sqrt{C_{12}^2(1 - (\tau + C_{12}^2))}} \\ + \sqrt{1 + \tau - 2\sqrt{C_{12}^2(1 - (\tau + C_{12}^2))}}\}$$

Thus we get the required bound Eq.(22). For MS subclass, $\lambda_0 = \frac{1}{\sqrt{2}}$, $\lambda_3 = \frac{\cos(\eta)}{\sqrt{2}}$ and $\lambda_4 = \frac{\sin(\eta)}{\sqrt{2}}$. So $\tau = \sin^2(\eta)$ and $C_{12}^2 = \cos^2(\eta)$. Hence the bound becomes:

$$1 + \sqrt{1 + \sin^2(\eta) + \cos^2(\eta)(1 - \sin^2(\eta) - \cos^2(\eta))} \\ + \sqrt{1 + \sin^2(\eta) - \cos^2(\eta)(1 - \sin^2(\eta) - \cos^2(\eta))} \\ = 1 + 2\sqrt{1 + \sin^2(\eta)}$$

which is the bound (20) for Maximal Slice states.

X. APPENDIX C

Calculation of monogamy score δ_D of GGHZ class and subclass S of extended GHZ class of states: The mutual information, for a quantum state ρ_{AB} , is given by [81–85]:

$$\mathcal{I}(\rho_{AB}) = \mathcal{S}(\rho_A) + \mathcal{S}(\rho_B) - \mathcal{S}(\rho_{AB}). \quad (42)$$

where $\mathcal{S}(\rho)$ denotes the Von Neumann entropy of the state ρ . The classical mutual information for a bipartite quantum state ρ_{AB} :

$$\mathcal{J}(\rho_{AB}) = \mathcal{S}(\rho_A) - \mathcal{S}(\rho_{A/B}). \quad (43)$$

$\mathcal{S}(\rho_{A/B})$ is the quantum conditional entropy given by:

$$\mathcal{S}(\rho_{A/B}) = \min_{\Pi_i^B} \sum_i p_i \mathcal{S}(\rho_{A/i}) \quad (44)$$

where the minimization is over all rank-1 measurements, Π_i^B performed over subsystem B . Here p_i is the corresponding probability for obtaining the outcome i and $\rho_{A/i}$ denotes the corresponding post measurement state of the subsystem A . The quantum discord [62, 63] of a state ρ_{AB} is given by:

$$\mathcal{D}(\rho_{AB}) = \mathcal{I}(\rho_{AB}) - \mathcal{J}(\rho_{AB}). \quad (45)$$

For GGHZ class of states, $\rho_{ABC} = |\varphi_{GGHZ}\rangle\langle\varphi_{GGHZ}|$. $\rho_{AB} = \rho_{AC} = \cos^2(\eta)|00\rangle\langle 00| + \sin^2(\eta)|11\rangle\langle 11|$. Now both of ρ_{AB} and ρ_{AC} are classical quantum states (i.e., they are of the form: $\sum_{p_k} \rho_k \otimes |k\rangle\langle k|$). Hence $\mathcal{D}(\rho_{AB}) = \mathcal{D}(\rho_{AC}) = 0$ (discord of a classical state vanishes). $\rho_{A:BC}$ is a bipartite (with A as one party and B and C together taken as another party) pure state. As quantum discord of a bipartite pure state is equal to its measure of entanglement, $\mathcal{D}(\rho_{A:BC})$ is given by the Von Neumann entropy of the corresponding reduced states. Hence

$$\mathcal{D}(\rho_{A:BC}) = \mathcal{S}(\rho_A)$$

$$= -(\cos^2 \eta \log_2(\cos^2 \eta) + \sin^2 \eta \log_2(\sin^2 \eta)).$$

This gives the the discord monogamy score of $|\varphi_{GHZ}\rangle$. For subclass S of extended GHZ class of states $\rho_{ABC} = |\chi\rangle\langle\chi|$.

$$\rho_{AB} = \lambda_0^2|00\rangle\langle 00| + (1 - \lambda_0^2)|11\rangle\langle 11| + \lambda_3\lambda_0(|00\rangle\langle 11| + |11\rangle\langle 00|).$$

In order to calculate the discord of a state one has to calculate Eq.(44) which is hard to compute for obvious reasons. However ρ_{AB} belongs to the class of X states for which an algorithm calculating discord is reported in [86]. Following the technique used there we get

$$\frac{-\ln 4(\lambda_0^2 \ln(\lambda_0^2) + (1 - \lambda_0^2) \ln(1 - \lambda_0^2)) + \ln(2)(1 + r \log_2(\frac{1+r}{2}) + 1 - r \log_2(\frac{1-r}{2}))}{\ln(2) \ln(4)}$$

where $r = \sqrt{1 + 4\lambda_0^4 + 4\lambda_0^2(-1 + \lambda_3^2)}$.

Now $\rho_{AC} = \lambda_0^2|00\rangle\langle 00| + \lambda_4^2|11\rangle\langle 11| + \lambda_3^2|10\rangle\langle 10| + T_3(|11\rangle\langle 10| + |10\rangle\langle 11|)$. It is a classical quantum state of the form $|k\rangle\langle k| \otimes \sum_{p_k} \rho_k$. Hence $D(\rho_{AC}) = 0$. Here $\rho_{A:BC}$ is a bipartite (with A as one party and B and C together taken as another party) pure state. Hence $D(\rho_{A:BC}) = -(\lambda_0^2 \log_2(\lambda_0^2) + (1 - \lambda_0^2) \log_2(1 - \lambda_0^2))$. On further simplification and using $\tau = 4\lambda_0^2\lambda_4^2$ and $C_{12}^2 = 4\lambda_0^2\lambda_3^2$ in Eq.(12), we get the discord monogamy score (23).

XI. APPENDIX D

Relation between τ and C_{12}^2 for states belonging to extended GHZ class of states characterized by $\lambda_0 = \frac{1}{\sqrt{2}}$: As $\lambda_0 = \frac{1}{\sqrt{2}}$,

$$\tau = 2\lambda_4^2 \quad (46)$$

and

$$C_{12}^2 = 2\lambda_3^2 \quad (47)$$

Hence

$$\tau = 1 - C_{12}^2. \quad (48)$$

Hence $\tau > \frac{1-C_{12}^2}{2}$ which in turn implies that for $\lambda_0 = \frac{1}{\sqrt{2}}$ Svetlichny inequality(2) is always violated. Now we focus on the range of τ occurring in the bound of the 99th facet inequality(6). We claim that τ always lies inside the interval $(\frac{C_{12}^2(1-C_{12}^2)}{1+C_{12}^2}, 1]$. If possible let τ lies outside the interval $(\frac{C_{12}^2(1-C_{12}^2)}{1+C_{12}^2}, 1]$, i.e. there exists some value τ lying outside this interval. This gives $\tau < \frac{C_{12}^2(1-C_{12}^2)}{1+C_{12}^2}$ which along with Eqs.(46, 47) implies $1 < 0$. This proves our claim which in turn implies for $\lambda_0 = \frac{1}{\sqrt{2}}$, 99 facet inequality(6) is violated for any positive value of τ and hence by any tripartite pure state belonging to the subclass S and characterized by $\lambda_0 = \frac{1}{\sqrt{2}}$.