

Quark mixing in an S_3 symmetric model with two Higgs doublets

Dipankar Das,^{a,*} Ujjal Kumar Dey,^{b,†} Palash B. Pal^{c,‡}

^a*Department of Physics, University of Calcutta, 92 Acharya Prafulla Chandra Road, Kolkata 700009, India*

^b*Centre for Theoretical Studies, Indian Institute of Technology Kharagpur, Kharagpur 721302, India*

^c*Theory Division, Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700064, India*

Abstract

We construct a model where the smallness of the masses of first quark generations implies the near block diagonal nature of the CKM matrix and vice-versa. For this set-up, we rely on a 2HDM structure with an S_3 symmetry. We show that an SM-like Higgs emerges naturally from such a construction. Moreover, the ratio of two VEVs, $\tan\beta$ can be precisely determined from the requirement of the near masslessness of the up- and down-quarks. The FCNC structure that arises from our model is also very predictive.

The Standard Model (SM) does not provide any connection between quark masses and mixings: they are independent parameters to be fixed by the experimental observations. One attractive way to obtain insight into these parameters is to impose some additional symmetry under which the generations of quarks transform in a non-trivial way. There have been many attempts where, by imposing a discrete symmetry on different generations of SM fermions, some relations between the masses and mixings have been obtained (see [1, 2] for review).

In this article, we present an attempt to relate two features of quark masses and mixings. The first of these two is the fact that the first generation quarks are very light compared to the other ones whereas the second concerns the near block-diagonal structure of the quark mixing matrix, or the Cabibbo-Kobayashi-Maskawa (CKM) matrix. This second feature comes out very clearly in the Wolfenstein parametrization [3] of the CKM matrix, where each element is written in a power series of a small parameter λ . If we keep only terms up to the linear order in λ , the CKM matrix is indeed block-diagonal. We propose a connection between these two features by invoking an S_3 symmetry.

Many works on flavour model building using S_3 symmetry have been done in the past [4–26]. In these constructions one usually employs, for the scalar sector, a three Higgs doublet structure which goes well with the aesthetic idea of having three replicas of Higgs doublets in conformity with three generations of fermions [27–41]. Even more complicated scalar structures are not uncommon [42–48]. However, in this paper, we rely on a two Higgs-doublet model (2HDM) scalar structure [49] which is much more economical in terms of independent parameters. Although the idea of a 2HDM with S_3 symmetry has been conceived lately [24], some distinct implications have not been emphasised earlier. For example, we will show that an S_3 symmetric 2HDM potential naturally delivers an SM-like Higgs boson which can be identified with the scalar resonance observed at the LHC with signal strengths

*ddphy@caluniv.ac.in

†ujjal@cts.iitkgp.ernet.in

‡pupal@theory.saha.ernet.in

in close agreement with the SM predictions [50]. We will also demonstrate how, in our scenario, the requirement of near-masslessness for the first generation of quarks dictates a particular value of $\tan\beta$, which will simultaneously render the CKM matrix block-diagonal. For intuitive understanding of the model Lagrangian and the conclusions that follow from it, a brief overview of the S_3 symmetry is in order.

The discrete symmetry group S_3 has three irreducible representations: $\mathbf{1}$, $\mathbf{1}'$ and $\mathbf{2}$. We pick a basis such that the generators in the $\mathbf{2}$ representation are given by

$$a = \begin{bmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}, \quad b = \begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}. \quad (1)$$

Note that a is of order 3, whereas b is of order 2. The rest of the elements can be obtained by taking products of powers of these two elements. In this basis the quark fields transform under S_3 in the following way:

$$\mathbf{2} : \begin{bmatrix} Q_1 \\ Q_2 \end{bmatrix}, \begin{bmatrix} u_{1R} \\ u_{2R} \end{bmatrix}, \begin{bmatrix} d_{1R} \\ d_{2R} \end{bmatrix}, \quad (2a)$$

$$\mathbf{1} : Q_3, u_{3R}, d_{3R}, \quad (2b)$$

where the Q_A 's ($A = 1, 2, 3$) are the usual left-handed SU(2) quark doublets, whereas the u_{AR} 's and d_{AR} 's are the right-handed up-type and down-type quark fields respectively, which are singlets of the SU(2) part of the gauge symmetry. Note that the square brackets, in Eqs. (1) and (2) as well as in the subsequent text, denote the doublet representation of S_3 , and has nothing to do with the representation of the enclosed fields under SU(2). Similarly, in the Higgs sector, there are two SU(2) doublets ϕ_i ($i = 1, 2$), and their transformation under the S_3 symmetry is as follows:

$$\mathbf{2} : \begin{bmatrix} \phi_1 \\ \phi_2 \end{bmatrix} \equiv \Phi. \quad (3)$$

We write the potential of the theory as follows:

$$V(\Phi) = V_2(\Phi) + V_4(\Phi), \quad (4a)$$

where $V_2(\Phi)$ contains only quadratic terms in the fields ϕ_1 and ϕ_2 whereas $V_4(\Phi)$ contains quartic terms. The most general S_3 -invariant form for $V_4(\Phi)$ is:

$$V_4(\Phi) = \lambda_1(\phi_1^\dagger\phi_1 + \phi_2^\dagger\phi_2)^2 + \lambda_2(\phi_1^\dagger\phi_2 - \phi_2^\dagger\phi_1)^2 + \lambda_3 \left\{ (\phi_1^\dagger\phi_2 + \phi_2^\dagger\phi_1)^2 + (\phi_1^\dagger\phi_1 - \phi_2^\dagger\phi_2)^2 \right\}. \quad (4b)$$

For the quadratic terms, the most general form is given by

$$V_2(\Phi) = \mu_1^2(\phi_1^\dagger\phi_1) + \mu_2^2(\phi_2^\dagger\phi_2) - \left(\mu_{12}^2\phi_1^\dagger\phi_2 + \text{h.c.} \right), \quad (4c)$$

which is not S_3 -symmetric unless the co-efficients satisfy some special conditions. If these conditions are not met, $V_2(\Phi)$ contains terms which softly break the S_3 symmetry, and we allow for such terms. We will consider various scenarios with the quadratic terms in a short while.

The parameters in the quartic part of the potential must be real because of hermiticity of the Lagrangian. In the quadratic part $V_2(\Phi)$, the parameters μ_1^2 and μ_2^2 are also real. The parameter μ_{12}^2 can be complex, but its phase can be absorbed by redefining either ϕ_1 or ϕ_2 . Thus, all parameters in $V(\Phi)$ can be taken to be real without any loss of generality. It has been argued [32] that in this case

the vacuum expectation values (VEVs) can also be taken to be real. Denoting the VEV of ϕ_i by v_i we write the doublets after symmetry breaking in the form

$$\phi_i = \begin{pmatrix} \phi_i^+ \\ \frac{1}{\sqrt{2}}(v_i + h_i + i\zeta_i) \end{pmatrix}, \quad (5)$$

and use the standard notation

$$v_1 = v \cos \beta, \quad v_2 = v \sin \beta, \quad (6)$$

where the W and Z -boson masses are proportional to $v \approx 246$ GeV. Assuming both v_1 and v_2 to be non-zero, the minimisation condition of the potential $V(\Phi)$ can be written as

$$\mu_1^2 = \mu_{12}^2 \tan \beta - (\lambda_1 + \lambda_3)v^2, \quad (7a)$$

$$\mu_2^2 = \mu_{12}^2 \cot \beta - (\lambda_1 + \lambda_3)v^2. \quad (7b)$$

Let us discuss the physical scalar spectrum of the model. We begin with the charged boson sector. One combination of ϕ_1^\pm and ϕ_2^\pm , to be denoted by w^\pm , will constitute an unphysical field that does not appear in the physical spectrum. The orthogonal combination, H^\pm , will be a physical charged scalar. The two combinations will be given by

$$\begin{pmatrix} w^\pm \\ H^\pm \end{pmatrix} = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix} \begin{pmatrix} \phi_1^\pm \\ \phi_2^\pm \end{pmatrix}. \quad (8)$$

The mass of the physical charged scalar can be easily calculated:

$$M_{H^\pm}^2 = \frac{2\mu_{12}^2}{\sin 2\beta} - 2\lambda_3 v^2. \quad (9)$$

In the pseudoscalar sector, there is one combination, z , which becomes unphysical after symmetry breaking, and there is one physical pseudoscalar field A . They are given by

$$\begin{pmatrix} z \\ A \end{pmatrix} = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix} \begin{pmatrix} \zeta_1 \\ \zeta_2 \end{pmatrix} \quad (10)$$

with

$$M_A^2 = \frac{2\mu_{12}^2}{\sin 2\beta} - 2(\lambda_2 + \lambda_3)v^2. \quad (11)$$

The mass matrix for the scalar part can be written as,

$$V_{\text{mass}}^S = (h_1 \quad h_2) \frac{1}{2} \mathbb{M}_S^2 \begin{pmatrix} h_1 \\ h_2 \end{pmatrix} \quad (12)$$

with

$$\mathbb{M}_S^2 = \begin{pmatrix} \mu_{12}^2 \frac{v_2}{v_1} + 2v_1^2(\lambda_1 + \lambda_3) & -\mu_{12}^2 + 2v_1 v_2(\lambda_1 + \lambda_3) \\ -\mu_{12}^2 + 2v_1 v_2(\lambda_1 + \lambda_3) & \mu_{12}^2 \frac{v_1}{v_2} + 2v_2^2(\lambda_1 + \lambda_3) \end{pmatrix}. \quad (13)$$

The diagonalisation of \mathbb{M}_S^2 will lead to two neutral physical scalars H and h ,

$$\begin{pmatrix} h \\ H \end{pmatrix} = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix} \begin{pmatrix} h_1 \\ h_2 \end{pmatrix} \quad (14)$$

with

$$m_H^2 = \frac{2\mu_{12}^2}{\sin 2\beta}, \quad m_h^2 = 2(\lambda_1 + \lambda_3)v^2. \quad (15)$$

At this point one should note that in the case of 2HDMs, the combination $H^0 = (v_1 h_1 + v_2 h_2)/v$ has SM-like couplings at the tree level. But in general H^0 is not a physical eigenstate. In the limit where H^0 is aligned with one of the physical CP -even scalars, is known as the *alignment limit* for 2HDMs. Eq. (14) shows that this is indeed the case in the present model, viz., that the eigenstate h is the same as H^0 . Thus the alignment limit emerges naturally [51] in our scenario. Hence by identifying h with the 125 GeV scalar observed at the LHC, our model becomes consistent, by design, with the LHC Higgs data [50].

Looking at the spectrum, we can identify the following different scenarios in regard to Eq. (4c).

1. If $\mu_1^2 = \mu_2^2$ and $\mu_{12}^2 = 0$, $V_2(\Phi)$ is completely S_3 -symmetric. In fact, the potential is invariant under a much bigger symmetry: an $SO(2)$ symmetry under which

$$\begin{aligned} \phi_1 &\rightarrow \phi_1 \cos \alpha - \phi_2 \sin \alpha, \\ \phi_2 &\rightarrow \phi_1 \sin \alpha + \phi_2 \cos \alpha. \end{aligned} \quad (16)$$

Thus, after gauge symmetry breaking when the ϕ_i 's develop vacuum expectation values (VEVs), we will have a massless scalar, a Goldstone boson as seen clearly from Eq. (15). This is not the scenario that we advocate.

2. If $\mu_1^2 \neq \mu_2^2$ and $\mu_{12}^2 = 0$, the potential is not S_3 symmetric, but Eq. (15) shows that we will still have a massless boson. Thus, this is not our desired scenario either.
3. If $\mu_1^2 = \mu_2^2$ and $\mu_{12}^2 \neq 0$, there exists no massless scalar, but Eq. (7) shows that we will now have $\tan \beta = 1$ or $v_1 = v_2$ because the potential has an exchange symmetry $\phi_1 \leftrightarrow \phi_2$. As we discuss later, this scenario will be detrimental to our aim.
4. If $\mu_1^2 \neq \mu_2^2$ and $\mu_{12}^2 \neq 0$, there is no massless scalar and also $\tan \beta$ can be arbitrary. This is the scenario that will be useful for us, implying that the soft-breaking terms are absolutely necessary.

We now present the most general Yukawa couplings involving the u_R quarks that is consistent with the gauge and S_3 symmetries. The S_3 symmetry cuts down on the number of Yukawa couplings drastically, and we obtain only the following couplings involving right-chiral u -type quarks:

$$\begin{aligned} \mathcal{L}_Y^{(u)} &= -A_u (\overline{Q}_1 \tilde{\phi}_1 + \overline{Q}_2 \tilde{\phi}_2) u_{3R} - B_u \left\{ (\overline{Q}_1 \tilde{\phi}_2 + \overline{Q}_2 \tilde{\phi}_1) u_{1R} + (\overline{Q}_1 \tilde{\phi}_1 - \overline{Q}_2 \tilde{\phi}_2) u_{2R} \right\} \\ &\quad - C_u \overline{Q}_3 (\tilde{\phi}_1 u_{1R} + \tilde{\phi}_2 u_{2R}) + \text{h.c.} \end{aligned} \quad (17)$$

We have used the standard abbreviation $\tilde{\phi}_i = i\sigma_2 \phi_i^*$. The Yukawa couplings of the d_R quarks can be obtained by replacing u_{AR} by d_{AR} , $\{A, B, C\}_u$ by $\{A, B, C\}_d$, and $\tilde{\phi}_i$ by ϕ_i in Eq. (17). Although the Yukawa couplings, in general, may be complex, we will discuss later that all but one phase can be absorbed in the field redefinitions.

After symmetry breaking, the mass matrices that arise in the quark sector have the following form:

$$\mathcal{M}_q = \frac{v}{\sqrt{2}} \begin{pmatrix} B_q \sin \beta & B_q \cos \beta & A_q \cos \beta \\ B_q \cos \beta & -B_q \sin \beta & A_q \sin \beta \\ C_q \cos \beta & C_q \sin \beta & 0 \end{pmatrix}, \quad (18)$$

where the subscripted index q can take the value u for the up-type quarks, and d for the down-type quarks. It is well-known that these matrices can be diagonalized through bi-unitary transformations, e.g., one can find two unitary matrices U_u and V_u , for the up-sector, such that $U_u \mathcal{M}_u V_u^\dagger$ is diagonal. The CKM matrix is then given by $U_u U_d^\dagger$.

The matrices U_u and U_d are the unitary matrices which diagonalize, through similarity transformations, the hermitian matrices $\mathcal{M}_u \mathcal{M}_u^\dagger$ and $\mathcal{M}_d \mathcal{M}_d^\dagger$ respectively. From Eq. (18), we obtain

$$\mathcal{M}_q \mathcal{M}_q^\dagger = \frac{1}{2} v^2 \begin{pmatrix} a_q^2 \cos^2 \beta + b_q^2 & \frac{1}{2} a_q^2 \sin 2\beta & B_q C_q^* \sin 2\beta \\ \frac{1}{2} a_q^2 \sin 2\beta & a_q^2 \sin^2 \beta + b_q^2 & B_q C_q^* \cos 2\beta \\ B_q^* C_q \sin 2\beta & B_q^* C_q \cos 2\beta & c_q^2 \end{pmatrix}, \quad (19)$$

where $a_q = |A_q|$ etc. Clearly, the three eigenvalues of $\mathcal{M}_u \mathcal{M}_u^\dagger$ would be the mass squared of the three up sector quarks, namely m_u^2, m_c^2 and m_t^2 , and the three eigenvalues of $\mathcal{M}_d \mathcal{M}_d^\dagger$ would be m_d^2, m_s^2 and m_b^2 . The eigenvalues can be obtained by solving the characteristic equation of the general matrix in Eq. (19). Introducing the shorthand notation

$$x = \frac{2m^2}{v^2} \quad (20)$$

for any fermion with mass m , this characteristic equation has the following form:

$$x^3 - (a^2 + 2b^2 + c^2)x^2 + (a^2 + b^2)(b^2 + c^2)x - a^2 b^2 c^2 \sin^2 3\beta = 0, \quad (21)$$

with subscripts u or d attached to the Yukawa couplings, as the case may be. Note that this equation is free from the phase of BC^* , which is the only phase that is present in Eq. (19).

Looking at the Lagrangian of Eq. (17) and the corresponding Lagrangian involving d_{iR} , we see why only one phase is present in Eq. (19). Any phase of A_u and A_d can be absorbed by redefining the fields u_{3R} and d_{3R} . After this, both B_u and B_d can be made real by redefining the fields u_{1R}, u_{2R} and d_{1R}, d_{2R} . Finally, either C_u or C_d can be made real by choosing the phase of Q_3 , but one of them remains complex. Alternatively, one can make both C_u or C_d real first, by redefining the right-chiral quark fields, and then either B_u or B_d can be made real by choosing the phases of Q_1 and Q_2 . Either way, one of the C_q 's or one of the B_q 's can be complex in the most general case. In what follows, we will assume that all Yukawa couplings are real, and use the lower-case symbols for them.

Before entering into a discussion of the eigenvalues obtained as solutions of Eq. (21), let us have some idea of the form of the diagonalizing matrix. As a first step, we can diagonalize only the terms in Eq. (19) that are proportional to a_q^2 . This is done, e.g., by a matrix

$$\mathcal{U} = \begin{pmatrix} 0 & 0 & 1 \\ \sin \beta & -\cos \beta & 0 \\ \cos \beta & \sin \beta & 0 \end{pmatrix}. \quad (22)$$

Note that this matrix does not depend on the Yukawa couplings, and is therefore the same for the up-type and down-type mass matrices. Applying a similarity transformation with this matrix on $\mathcal{M} \mathcal{M}^\dagger$, we obtain

$$M^2 = \mathcal{U} \mathcal{M} \mathcal{M}^\dagger \mathcal{U}^\dagger = \frac{1}{2} v^2 \begin{pmatrix} c^2 & -bc \cos 3\beta & bc \sin 3\beta \\ -bc \cos 3\beta & b^2 & 0 \\ bc \sin 3\beta & 0 & a^2 + b^2 \end{pmatrix}, \quad (23)$$

with subscripts u and d attached for quarks of positive and negative charges respectively.

In the preamble of the article, we said that we want to relate the almost-masslessness of first generation quarks with the almost-block-diagonal form of the CKM matrix. We now narrow down the scenario in which we can have one zero eigenvalue in both up-type and down-type quark sector, as well as a block-diagonal CKM matrix.

First we note that if one solution of Eq. (21) is zero, then the x -independent term should vanish in that equation. In this case, the eigenvalues of $\mathcal{M}\mathcal{M}^\dagger$ are given by

$$0, \quad \frac{1}{2}v^2(b^2 + c^2), \quad \frac{1}{2}v^2(a^2 + b^2). \quad (24)$$

For the diagonalizing matrix, we now consider two different cases.

Case 1: Some Yukawa couplings vanish

Surely, the x -independent term in Eq. (21) can vanish if at least one of the Yukawa couplings is zero. Looking at Eq. (23), we see that $a = 0$ does not make M^2 block-diagonal, so we reject this possibility. If either b or c vanishes, the matrix M^2 becomes completely diagonal. This means that for b or $c = 0$, the same matrix \mathcal{U} will diagonalize both $\mathcal{M}_u\mathcal{M}_u^\dagger$ and $\mathcal{M}_d\mathcal{M}_d^\dagger$ making the CKM matrix a unit matrix. Therefore making some Yukawa coupling vanish to obtain one zero mass does not produce the desirable block-diagonal structure of the CKM matrix.

One should recall that making $\mu_1^2 = \mu_2^2$ in Eq. (4c) had led to $\tan\beta = 1$, which in view of Eq. (21) demands that one of the Yukawas must be zero in order to obtain zero mass eigenvalue. For this reason we discard this particular form of $V_2(\Phi)$.

Case 2: $\sin 3\beta = 0$

However, there is a second and more attractive possibility. From the characteristic equation, Eq. (21), one can see that zero eigenvalue can also be ensured if

$$\sin 3\beta = 0. \quad (25)$$

Discarding the trivial solution $\beta = 0$, we obtain the solution $\beta = \frac{1}{3}\pi$ which implies that $\tan\beta = \sqrt{3}$ i.e., $v_2 = \sqrt{3}v_1 = \sqrt{3}v/2$.¹ Looking at Eq. (23) now, we see that this value of β also makes the matrix M^2 block-diagonal, and one obtains

$$M^2 = \mathcal{U}\mathcal{M}\mathcal{M}^\dagger\mathcal{U}^\dagger = \frac{1}{2}v^2 \begin{pmatrix} c^2 & bc & 0 \\ bc & b^2 & 0 \\ 0 & 0 & a^2 + b^2 \end{pmatrix}. \quad (26)$$

Notice that the third generation has been singled out, and therefore $v\sqrt{(a^2 + b^2)}/2$ can be readily identified with the mass of the third generation quark. In order that it be much heavier than the quarks in the first two generations, we need

$$a^2 \gg b^2, c^2 \quad (27)$$

in both up and down sectors.

¹While this VEV alignment is useful for our consequent discussions we note that in case of a three Higgs doublet model with S_3 symmetry, the minimisation of potential leads to a vev alignment $v_1 = \sqrt{3}v_2$ and such alignment implies a residual \mathbb{Z}_2 symmetry [38]. In the present case, however, no such implications are possible.

Complete diagonalization would require a further similarity transformation affecting the upper 2×2 block. This will involve the values of the Yukawa couplings. Thus, we obtain

$$U_u = \mathcal{O}_u \mathcal{U}, \quad U_d = \mathcal{O}_d \mathcal{U}, \quad (28)$$

where

$$\mathcal{O}_q = \begin{pmatrix} \cos \theta_q & -\sin \theta_q & 0 \\ \sin \theta_q & \cos \theta_q & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (29)$$

with

$$\tan \theta_q = \frac{c_q}{b_q}. \quad (30)$$

From Eq. (28) the CKM matrix can now be written as,

$$V_{\text{CKM}} = U_u U_d^\dagger = \mathcal{O}_u \mathcal{O}_d^\dagger = \begin{pmatrix} \cos(\theta_u - \theta_d) & -\sin(\theta_u - \theta_d) & 0 \\ \sin(\theta_u - \theta_d) & \cos(\theta_u - \theta_d) & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (31)$$

Thus the difference $\theta_u - \theta_d$, which can be identified with the Cabibbo angle, θ_C .

In passing, we make a point about the VEV alignment, i.e., the value of β , dictated by Eq. (25). It reflects our choice of the representation for S_3 . Had we chosen a different representation, the value of β would in general be different. But the physical implications should be independent of the representation, and so the block-diagonal form of the CKM matrix would still result.

Having reproduced the leading order effects of the mixing matrix in the Wolfenstein parametrization as a consequence of the masslessness of the first generation quarks, we now explore whether one can do better. So far, the conclusions that we derived came from Eq. (25), which is a statement about the relative magnitude of the VEVs of the two Higgs doublets. Note that this relation is not protected by any symmetry. Suppose we deviate from Eq. (25) by a small amount such that

$$\sin 3\beta = \delta. \quad (32)$$

Since δ is expected to be small, we do not expect the heavier quark masses to be altered very much by this change. The sums of eigenvalues etc. will also not change appreciably. The only thing that will change dramatically is the product of all eigenvalues, which should be the x -independent term in Eq. (21). Therefore the first generation quark masses will be given, in the notation of Eq. (20), by

$$x_u = \frac{2m_u^2}{v^2} \approx \frac{a_u^2 b_u^2 c_u^2 \delta^2}{(a_u^2 + b_u^2)(b_u^2 + c_u^2)} \approx \frac{b_u^2 c_u^2 \delta^2}{b_u^2 + c_u^2} \quad (33a)$$

$$x_d = \frac{2m_d^2}{v^2} \approx \frac{a_d^2 b_d^2 c_d^2 \delta^2}{(a_d^2 + b_d^2)(b_d^2 + c_d^2)} \approx \frac{b_d^2 c_d^2 \delta^2}{b_d^2 + c_d^2}, \quad (33b)$$

where in the last step we have used the hierarchy mentioned in Eq. (27). Now, using Eq. (30), we can write

$$m_u^2 \approx \frac{1}{2} v^2 \delta^2 \times (b_u^2 + c_u^2) \sin^2 \theta_u \cos^2 \theta_u \approx \frac{1}{4} m_c^2 \delta^2 \sin^2 2\theta_u \quad (34a)$$

$$m_d^2 \approx \frac{1}{2} v^2 \delta^2 \times (b_d^2 + c_d^2) \sin^2 \theta_d \cos^2 \theta_d \approx \frac{1}{4} m_s^2 \delta^2 \sin^2 2\theta_d. \quad (34b)$$

Since $\theta_u - \theta_d = \theta_C$, Eqs. (34a) and (34b) can be solved for δ and θ_u or θ_d . Taking all the uncertainties into account we have found $\delta > 0.2$ which is inconsistent with our assumption of small δ in Eq. (32). Therefore, this minimal framework is not sufficient to reproduce the observed masses of the first generation quarks.

Now, for completeness, we comment on the flavour changing neutral currents (FCNC) in our model. To set up the notations we first lay out the Yukawa Lagrangian for 2HDM in the following form:

$$\mathcal{L}_Y = - \sum_{k=1}^2 \left[\bar{\mathbf{Q}}_L \Gamma_k \phi_k \mathbf{d}_R + \bar{\mathbf{Q}}_L \Delta_k \tilde{\phi}_k \mathbf{u}_R \right] + \text{h.c.}, \quad (35)$$

where we have kept the notation for the field the same as in Eq. (17) but put them in boldface font to remind ourselves that the generation indices have been suppressed. Unlike Eq. (17), here we also take into account the Yukawa Lagrangian for the down sector too. Here $\Delta_{1,2}$ and $\Gamma_{1,2}$ represent the Yukawa matrices in the up and down sectors respectively. By comparing Eqs. (35) and (17) we can write,

$$\Delta_1 = \begin{pmatrix} 0 & b_u & a_u \\ b_u & 0 & 0 \\ c_u & 0 & 0 \end{pmatrix}, \quad \Delta_2 = \begin{pmatrix} b_u & 0 & 0 \\ 0 & -b_u & a_u \\ 0 & c_u & 0 \end{pmatrix}, \quad (36)$$

and the Γ_k 's can be obtained by replacing the subscript u by the subscript d in the matrices. Now, the Yukawa Lagrangian in terms of physical fields can be written as

$$\begin{aligned} \mathcal{L}_{\text{Yuk}} = & -\frac{h}{v} (\bar{\mathbf{d}} D_d \mathbf{d} + \bar{\mathbf{u}} D_u \mathbf{u}) + \frac{H}{v} \left[\bar{\mathbf{d}} (N_d P_R + N_d^\dagger P_L) \mathbf{d} + \bar{\mathbf{u}} (N_u P_R + N_u^\dagger P_L) \mathbf{u} \right] \\ & -\frac{iA}{v} \left[\bar{\mathbf{u}} (N_u P_R - N_u^\dagger P_L) \mathbf{u} - \bar{\mathbf{d}} (N_d P_R - N_d^\dagger P_L) \mathbf{d} \right] \\ & + \frac{\sqrt{2}H^+}{v} \bar{\mathbf{u}} \left[V_{\text{CKM}} N_d P_R - N_u^\dagger V_{\text{CKM}} P_L \right] \mathbf{d} + \text{h.c.}, \end{aligned} \quad (37)$$

where D_u and D_d are the diagonal mass matrices in the up and down sectors respectively. Note that the SM-like scalar, h , does not have any FCNC couplings. This is a direct consequence of the natural alignment that we have talked about earlier. The matrices N_u and N_d , in Eq. (37), carry the information of FCNC in the up and down sectors respectively and are given by,

$$N_u = \frac{1}{\sqrt{2}} U_u (\Delta_1 v_2 - \Delta_2 v_1) V_u^\dagger, \quad (38a)$$

$$N_d = \frac{1}{\sqrt{2}} U_d (\Gamma_1 v_2 - \Gamma_2 v_1) V_d^\dagger. \quad (38b)$$

Note that the expressions for V_u and V_d can be obtained from diagonalizing $\mathcal{M}^\dagger \mathcal{M}$ for both up and down sectors. The matrices $\mathcal{M}^\dagger \mathcal{M}$ can be obtained from $\mathcal{M} \mathcal{M}^\dagger$ by making the interchange $a \leftrightarrow c$ in the Yukawa couplings. Because of this interchange, the matrix V is different from U in two respects. First, the matrix corresponding to \mathcal{U} should have the last two rows interchanged so that the eigenvalues can occur in the same order. Second, the angle θ_q should be replaced by θ'_q , which will be given by

$$\tan \theta'_q = \frac{a_q}{b_q}. \quad (39)$$

In view of the hierarchy mentioned in Eq. (27), we can use these to write

$$\sin \theta'_q \approx 1, \quad \cos \theta'_q \approx \frac{b_q}{a_q}, \quad (40)$$

neglecting higher order terms in b_q/a_q . Replacing the Yukawa couplings by the mass eigenvalues and the angles θ_q , we obtain

$$N_u \approx \begin{pmatrix} -\frac{3}{2}m_c \sin 2\theta_u & 0 & -m_t \sin \theta_u \\ \frac{1}{2}m_c(3 \cos^2 \theta_u - 1) & 0 & m_t \cos \theta_u \\ 0 & m_c \cos \theta_u & 0 \end{pmatrix}, \quad (41)$$

neglecting corrections of order m_c/m_t . A similar expression for N_d can be obtained from Eq. (41) by replacing θ_u, m_c, m_t with θ_d, m_s, m_b respectively. Thus the FCNCs are uniquely determined by θ_u or θ_d . One should keep in mind that this represents the FCNC couplings at the leading order, i.e., when the CKM matrix is block-diagonal. In a more complete framework where the CKM matrix can be reproduced exactly these FCNC matrices are expected to get small corrections.

A trivial but viable solution to the FCNC problem would be to make all the scalars except h sufficiently heavy. Moreover, the bounds from the electroweak T -parameter can also be evaded if the non-standard scalars, H, A and H^\pm are nearly degenerate [52, 53].

In summary, we connect two apparently disjoint experimental observations namely, the tiny masses of first generation of quarks and the near block-diagonal structure of the CKM matrix in a simple set-up of 2HDM with an S_3 symmetry. We attribute these two features of the quark sector to Nature's choice of a particular value of $\tan \beta$. An added bonus of our model is the existence of a light scalar, which can be identified with the 125 GeV Higgs observed at the LHC, in view of a naturally emerging alignment limit. Admittedly, the exact CKM matrix and correct non-zero masses for the first generation of quarks could not be reproduced in this minimalistic scenario. Perhaps our set-up can be taken as a constituent towards a more elaborate framework which can address the full quark structure.

Acknowledgements: DD thanks Arcadi Santamaria and Anirban Kundu for useful discussions. The work of UKD is supported by Department of Science and Technology, Government of India under the fellowship reference number PDF/2016/001087 (SERB National Post-Doctoral fellowship).

References

- [1] G. Altarelli and F. Feruglio, *Discrete Flavor Symmetries and Models of Neutrino Mixing*, *Rev. Mod. Phys.* **82** (2010) 2701–2729, [[arXiv:1002.0211](#)].
- [2] H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada, and M. Tanimoto, *Non-Abelian Discrete Symmetries in Particle Physics*, *Prog. Theor. Phys. Suppl.* **183** (2010) 1–163, [[arXiv:1003.3552](#)].
- [3] L. Wolfenstein, *Parametrization of the Kobayashi-Maskawa Matrix*, *Phys. Rev. Lett.* **51** (1983) 1945.
- [4] S. Pakvasa and H. Sugawara, *Discrete Symmetry and Cabibbo Angle*, *Phys. Lett.* **B73** (1978) 61–64.
- [5] Y. Koide, *Universal seesaw mass matrix model with an $S(3)$ symmetry*, *Phys.Rev.* **D60** (1999) 077301, [[hep-ph/9905416](#)].
- [6] J. Kubo, A. Mondragon, M. Mondragon, and E. Rodriguez-Jauregui, *The Flavor symmetry*, *Prog. Theor. Phys.* **109** (2003) 795–807, [[hep-ph/0302196](#)]. [Erratum: *Prog. Theor. Phys.*114,287(2005)].

- [7] P. Harrison and W. Scott, *Permutation symmetry, tri - bimaximal neutrino mixing and the S_3 group characters*, *Phys.Lett.* **B557** (2003) 76, [[hep-ph/0302025](#)].
- [8] S.-L. Chen, M. Frigerio, and E. Ma, *Large neutrino mixing and normal mass hierarchy: A Discrete understanding*, *Phys. Rev.* **D70** (2004) 073008, [[hep-ph/0404084](#)]. [Erratum: *Phys. Rev.* **D70**, 079905(2004)].
- [9] Y. Koide, *$S(3)$ symmetry and neutrino masses and mixings*, *Eur. Phys. J.* **C50** (2007) 809–816, [[hep-ph/0612058](#)].
- [10] A. Mondragon, M. Mondragon, and E. Peinado, *Lepton masses, mixings and FCNC in a minimal $S(3)$ -invariant extension of the Standard Model*, *Phys. Rev.* **D76** (2007) 076003, [[arXiv:0706.0354](#)].
- [11] C.-Y. Chen and L. Wolfenstein, *Consequences of approximate $S(3)$ symmetry of the neutrino mass matrix*, *Phys. Rev.* **D77** (2008) 093009, [[arXiv:0709.3767](#)].
- [12] R. Jora, J. Schechter, and M. Naeem Shahid, *Perturbed $S(3)$ neutrinos*, *Phys.Rev.* **D80** (2009) 093007, [[arXiv:0909.4414](#)].
- [13] T. Kaneko and H. Sugawara, *Broken S_3 Symmetry in Flavor Physics*, *Phys. Lett.* **B697** (2011) 329–332, [[arXiv:1011.5748](#)].
- [14] T. Teshima and Y. Okumura, *Quark/lepton mass and mixing in S_3 invariant model and CP-violation of neutrino*, *Phys. Rev.* **D84** (2011) 016003, [[arXiv:1103.6127](#)].
- [15] S. Dev, S. Gupta, and R. R. Gautam, *Broken S_3 Symmetry in the Neutrino Mass Matrix*, *Phys.Lett.* **B702** (2011) 28–33, [[arXiv:1106.3873](#)].
- [16] A. G. Dias, A. C. B. Machado, and C. C. Nishi, *An S_3 Model for Lepton Mass Matrices with Nearly Minimal Texture*, *Phys. Rev.* **D86** (2012) 093005, [[arXiv:1206.6362](#)].
- [17] D. Meloni, *S_3 as a flavour symmetry for quarks and leptons after the Daya Bay result on θ_{13}* , *JHEP* **05** (2012) 124, [[arXiv:1203.3126](#)].
- [18] S. Dev, R. R. Gautam, and L. Singh, *Broken S_3 Symmetry in the Neutrino Mass Matrix and Non-Zero θ_{13}* , *Phys. Lett.* **B708** (2012) 284–289, [[arXiv:1201.3755](#)].
- [19] S. Zhou, *Relatively large θ_{13} and nearly maximal θ_{23} from the approximate S_3 symmetry of lepton mass matrices*, *Phys. Lett.* **B704** (2011) 291–295, [[arXiv:1106.4808](#)].
- [20] E. Ma and B. Melic, *Updated S_3 model of quarks*, *Phys. Lett.* **B725** (2013) 402–406, [[arXiv:1303.6928](#)].
- [21] H. B. Benaoum, *Broken S_3 Neutrinos*, *Phys. Rev.* **D87** (2013) 073010, [[arXiv:1302.0950](#)].
- [22] E. Ma and R. Srivastava, *Dirac or inverse seesaw neutrino masses with $B - L$ gauge symmetry and S_3 flavor symmetry*, *Phys. Lett.* **B741** (2015) 217–222, [[arXiv:1411.5042](#)].
- [23] D. Das, U. K. Dey, and P. B. Pal, *S_3 symmetry and the quark mixing matrix*, *Phys. Lett.* **B753** (2016) 315–318, [[arXiv:1507.06509](#)].
- [24] D. Cogollo and J. P. Silva, *Two Higgs doublet models with an S_3 symmetry*, *Phys. Rev.* **D93** (2016), no. 9 095024, [[arXiv:1601.02659](#)].

- [25] S. Pramanick and A. Raychaudhuri, *Neutrino mass model with S_3 symmetry and seesaw interplay*, *Phys. Rev.* **D94** (2016), no. 11 115028, [[arXiv:1609.06103](#)].
- [26] I. de Medeiros Varzielas, R. W. Rasmussen, and J. Talbert, *Bottom-Up Discrete Symmetries for Cabibbo Mixing*, *Int. J. Mod. Phys.* **A32** (2017), no. 06n07 1750047, [[arXiv:1605.03581](#)].
- [27] N. G. Deshpande, M. Gupta, and P. B. Pal, *Flavor changing processes and CP violation in $S(3) \times Z(3)$ model*, *Phys. Rev.* **D45** (1992) 953–957.
- [28] J. Kubo, H. Okada, and F. Sakamaki, *Higgs potential in minimal $S(3)$ invariant extension of the standard model*, *Phys. Rev.* **D70** (2004) 036007, [[hep-ph/0402089](#)].
- [29] T. Teshima, *Flavor mass and mixing and $S(3)$ symmetry: An $S(3)$ invariant model reasonable to all*, *Phys. Rev.* **D73** (2006) 045019, [[hep-ph/0509094](#)].
- [30] Y. Koide, *Permutation symmetry $S(3)$ and VEV structure of flavor-triplet Higgs scalars*, *Phys. Rev.* **D73** (2006) 057901, [[hep-ph/0509214](#)].
- [31] O. F. Beltran, M. Mondragon, and E. Rodriguez-Jauregui, *Conditions for vacuum stability in an $S(3)$ extension of the standard model*, *J.Phys.Conf.Ser.* **171** (2009) 012028.
- [32] P. M. Ferreira, M. Maniatis, O. Nachtmann, and J. P. Silva, *CP properties of symmetry-constrained two-Higgs-doublet models*, *JHEP* **08** (2010) 125, [[arXiv:1004.3207](#)].
- [33] G. Bhattacharyya, P. Leser, and H. Pas, *Exotic Higgs boson decay modes as a harbinger of S_3 flavor symmetry*, *Phys. Rev.* **D83** (2011) 011701, [[arXiv:1006.5597](#)].
- [34] G. Bhattacharyya, P. Leser, and H. Pas, *Novel signatures of the Higgs sector from S_3 flavor symmetry*, *Phys. Rev.* **D86** (2012) 036009, [[arXiv:1206.4202](#)].
- [35] T. Teshima, *Higgs potential in S_3 invariant model for quark/lepton mass and mixing*, *Phys. Rev.* **D85** (2012) 105013, [[arXiv:1202.4528](#)].
- [36] A. C. B. Machado and V. Pleitez, *A model with two inert scalar doublets*, *Annals Phys.* **364** (2016) 53–67, [[arXiv:1205.0995](#)].
- [37] E. Barradas-Guevara, O. Félix-Beltrán, and E. Rodríguez-Jáuregui, *Trilinear self-couplings in an $S(3)$ flavored Higgs model*, *Phys. Rev.* **D90** (2014), no. 9 095001, [[arXiv:1402.2244](#)].
- [38] D. Das and U. K. Dey, *Analysis of an extended scalar sector with S_3 symmetry*, *Phys. Rev.* **D89** (2014), no. 9 095025, [[arXiv:1404.2491](#)]. [Erratum: *Phys. Rev.* D91,no.3,039905(2015)].
- [39] E. Barradas-Guevara, O. Félix-Beltrán, and E. Rodríguez-Jáuregui, *CP breaking in $S(3)$ flavoured Higgs model*, [arXiv:1507.05180](#).
- [40] M. Merchand and M. Sher, *Three doublet lepton-specific model*, *Phys. Rev.* **D95** (2017), no. 5 055004, [[arXiv:1611.06887](#)].
- [41] D. Emmanuel-Costa, O. M. Ogreid, P. Osland, and M. N. Rebelo, *Spontaneous symmetry breaking in the S_3 -symmetric scalar sector*, *JHEP* **02** (2016) 154, [[arXiv:1601.04654](#)]. [Erratum: *JHEP*08,169(2016)].
- [42] Z.-z. Xing, D. Yang, and S. Zhou, *Broken S_3 Flavor Symmetry of Leptons and Quarks: Mass Spectra and Flavor Mixing Patterns*, *Phys. Lett.* **B690** (2010) 304–310, [[arXiv:1004.4234](#)].

- [43] F. Gonzalez Canales, A. Mondragon, and M. Mondragon, *The S_3 Flavour Symmetry: Neutrino Masses and Mixings*, *Fortsch. Phys.* **61** (2013) 546–570, [[arXiv:1205.4755](#)].
- [44] F. González Canales, A. Mondragón, M. Mondragón, U. J. Saldaña Salazar, and L. Velasco-Sevilla, *Quark sector of S_3 models: classification and comparison with experimental data*, *Phys. Rev.* **D88** (2013) 096004, [[arXiv:1304.6644](#)].
- [45] A. E. Cárcamo Hernández, R. Martinez, and J. Nisperuza, *S_3 discrete group as a source of the quark mass and mixing pattern in 331 models*, *Eur. Phys. J.* **C75** (2015), no. 2 72, [[arXiv:1401.0937](#)].
- [46] A. E. Cárcamo Hernández, I. de Medeiros Varzielas, and E. Schumacher, *Fermion and scalar phenomenology of a two-Higgs-doublet model with S_3* , *Phys. Rev.* **D93** (2016), no. 1 016003, [[arXiv:1509.02083](#)].
- [47] A. E. Cárcamo Hernández, I. de Medeiros Varzielas, and N. A. Neill, *Novel Randall-Sundrum model with S_3 flavor symmetry*, *Phys. Rev.* **D94** (2016), no. 3 033011, [[arXiv:1511.07420](#)].
- [48] J. C. Gómez-Izquierdo, *Non-Minimal Flavored $S_3 \otimes Z_2$ Left-Right Symmetric Model*, [[arXiv:1701.01747](#)].
- [49] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, and J. P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1–102, [[arXiv:1106.0034](#)].
- [50] **ATLAS, CMS** Collaboration, G. Aad et al., *Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV*, *JHEP* **08** (2016) 045, [[arXiv:1606.02266](#)].
- [51] P. S. Bhupal Dev and A. Pilaftsis, *Maximally Symmetric Two Higgs Doublet Model with Natural Standard Model Alignment*, *JHEP* **12** (2014) 024, [[arXiv:1408.3405](#)]. [Erratum: *JHEP*11,147(2015)].
- [52] W. Grimus, L. Lavoura, O. M. Ogreid, and P. Osland, *A Precision constraint on multi-Higgs-doublet models*, *J. Phys.* **G35** (2008) 075001, [[arXiv:0711.4022](#)].
- [53] G. Bhattacharyya, D. Das, P. B. Pal, and M. N. Rebelo, *Scalar sector properties of two-Higgs-doublet models with a global $U(1)$ symmetry*, *JHEP* **10** (2013) 081, [[arXiv:1308.4297](#)].