

Effect of smoking on semen quality, FSH, testosterone level, and CAG repeat length in androgen receptor gene of infertile men in an Indian city

Amrita Mitra, Baidyanath Chakraborty, Dyutiman Mukhopadhyay, Manisha Pal, Sanjit Mukherjee, Samir Banerjee & Keya Chaudhuri

To cite this article: Amrita Mitra, Baidyanath Chakraborty, Dyutiman Mukhopadhyay, Manisha Pal, Sanjit Mukherjee, Samir Banerjee & Keya Chaudhuri (2012) Effect of smoking on semen quality, FSH, testosterone level, and CAG repeat length in androgen receptor gene of infertile men in an Indian city, *Systems Biology in Reproductive Medicine*, 58:5, 255-262, DOI: [10.3109/19396368.2012.684195](https://doi.org/10.3109/19396368.2012.684195)

To link to this article: <https://doi.org/10.3109/19396368.2012.684195>



Published online: 14 May 2012.



Submit your article to this journal [↗](#)



Article views: 1304



View related articles [↗](#)



Citing articles: 18 View citing articles [↗](#)

CLINICAL CORNER: COMMUNICATION

Effect of smoking on semen quality, FSH, testosterone level, and CAG repeat length in androgen receptor gene of infertile men in an Indian city

Amrita Mitra¹, Baidyanath Chakraborty¹, Dyutiman Mukhopadhyay², Manisha Pal³, Sanjit Mukherjee⁴, Samir Banerjee², and Keya Chaudhuri⁴

¹Institute of Reproductive Medicine, Kolkata, ²Department of Zoology, University of Calcutta, Kolkata, ³Department of Statistics, University of Calcutta, Kolkata, ⁴Molecular and Human Genetics Division, Indian Institute of Chemical Biology, Kolkata, India

This study was conducted as part of an epidemiological survey of 126 nonsmokers and 178 smokers, showing primary infertility residing around Kolkata region of Eastern India. Their life-style history including smoking habits along with semen and blood were collected. The study examined the association of cigarette smoking with the risk of infertility, by determining the semen quality, follicle stimulating hormone (FSH), luteinizing hormone (LH), testosterone levels, and androgen receptor (AR)-CAG repeat length in a group of smokers compared with a control group (non smokers). Based on conventional WHO criteria, lower sperm motility ($P < 0.001$) and increased sperm morphological defects ($P < 0.0001$) were associated with smoking habits. Binary logistic regression analysis for the effect of smoking status on sperm DNA integrity demonstrated significant positive correlation ($p = 0.006$). Serum FSH and LH levels were higher for smokers compared with non-smokers while the testosterone level decreased significantly with the increasing smoking habit. The mean length of CAG repeats in AR gene was significantly higher for smokers with low testosterone compared to non-smokers. The study suggested that smoking is associated with altered semen quality, endocrine hormonal status, and number of CAG repeats in the AR gene.

Keywords CAG repeats, DNA integrity, reproductive hormone, semen quality, smoking

Introduction

Infertility is traditionally defined as the inability of a couple to conceive following one year of unprotected intercourse. It is not an uncommon problem; as the numbers of infertile couples are rising day by day. In those couples who seek evaluation from a reproductive specialist, more than 30% of males have defective sperm quality that significantly contributes to their infertility [Gaur et al. 2007]. There are several factors which contribute to the increase in the male factor infertility [Lo et al. 2005]. Among these, exposure to

several environmental, occupational, and behavioral factors have been reported as risk factors for male infertility. Since cigarette smoke contains mutagens and carcinogens and smoking is very common all over the world, there have been increased concerns of the unfavorable effects of smoking on male reproductive function [Mukhopadhyay et al. 2010].

Semen analysis is frequently used to assess male infertility in clinical andrology. Human sperm quality is usually defined by standard WHO semen analysis parameters: count, motility, and morphology [WHO 1999]. Recently the relationship between DNA integrity and conventional semen parameters has been established. An increased percentage of DNA damage is associated with a rise in semen abnormality [Trisini et al. 2004; Varghese et al. 2009].

Environmental factors can affect the male reproductive system either directly, causing decreased or altered sperm production in the testis, and/or indirectly through the endocrine system causing hormonal imbalance. The levels of follicle-stimulating hormone (FSH), luteinizing hormone (LH), and testosterone could be correlated with male infertility [Padron et al. 1989]. Androgen is an important steroid hormone responsible for normal spermatogenesis, decreasing the level of this hormone leads to impaired sperm production. All androgens act through a single intracellular androgen receptor (AR), which is encoded by a single-copy gene mapped on the long arm (Xq11-12) of the X-chromosome [Chang et al. 1988]. The AR gene has 8 exons, of which exon 1 harbors a CAG repeat motif coding for a polyglutamine stretch. Reduced androgenicity in 20% of infertile men is caused by the expansion of CAG tract [Yong et al. 1998].

The present study undertook an analysis of semen parameters such as sperm count, motility, morphology, DNA integrity, endocrine hormones, and CAG repeats in the AR gene among smoker and non-smoker infertile men to assess the association of cigarette smoking with infertility in this population of the eastern region of India.

Received 22 September 2011; accepted 11 February 2012.

*Address correspondence to Dr. Keya Chaudhuri, Chief Scientist, Molecular & Human Genetics Division and Head, Academic Affairs Division Indian Institute of Chemical Biology, 4 Raja S C Mullick Road, Kolkata-700032, India. Email: kchaudhuri@iicb.res.in or keya.chaudhuri@gmail.com

Results and Discussion

The study population consisted of patients attending our infertility clinic for initial evaluation. Patients of age group 30-60 y were included in the study after a thorough investigation of their past medical history and habits. The body mass index (BMI) for all of the patients was calculated from the information obtained, nonsmokers and light smokers showed almost similar BMI (25.92 ± 4.77 and 25.68 ± 5.92 , respectively); in comparison, the mean BMI among heavy smokers were significantly lower being 21.31 ± 4.22 kg/m² (Table 1). The sperm count among the heavy smokers ($50 \times 10^6 \pm 20$) and light smokers ($76 \times 10^6 \pm 50$) was significantly lower compared to non-smokers ($130 \times 10^6 \pm 50$, $p = 0.001$). Among the nonsmokers, most of the patients were normozoospermic (60.3%), other seminal phenotypes are also present: teratozoospermic (19%), asthenozoospermic (4%), and about 24.1% showed sperm DNA damage. Most of the light smokers presented with normozoospermic (50%), azoospermic (14.3%), and teratozoospermic (16.3%) phenotype, with 42.5% of the patients showing damaged sperm DNA integrity. Heavy smokers produced strikingly different parameters compared to the other two groups. About 85% of the patients showed a defect in seminal phenotype; patients were mostly azoospermic (22.5%), oligozoospermic (20%), and teratozoospermic (30%); about 37.9% presented head defects and 43% sperm DNA damage was prevalent among the heavy smokers (Table 2).

The frequency χ^2 -test shows significant association between smoking habit and seminal phenotype ($\chi^2 = 59.462$, p -value ≈ 0.000). In order to begin to understand the nature of association, logistic regression analysis was carried out as smoking habit and seminal phenotype are categorical variables, as follows: phenotype normozoospermic (0, Reference Event), asthenozoospermic (5), teratozoospermic (4), severe oligozoospermic (3), oligozoospermic (2), and finally azoospermic (1) (Table 3A). A borderline effect of BMI upon seminal phenotype was noted (OR 0.87 95% CI (0.75-1.00), $p = 0.05$) in logit 1, with an indication for the effect of BMI upon decreasing phenotype which tends to shift from normo- to asthenozoospermia. In logit 5 the BMI is significantly lower in the azoospermic group ($p < 0.001$). Logits 1, 2, 3, and 5 indicate a phenotypical changes from normozoospermia to azo/oligo/severe oligo/asthenozoospermia respectively with increasing smoking habits ($p < 0.05$) (Table 3A).

Table 2. Characteristics of semen, reproductive hormones for participants according to level of current smoking.

Seminal characteristics	Non-smoker	Light smokers (1-20 cigarettes/day)	Heavy smokers (>20 cigarettes/day)
Seminal phenotype			
Normozoospermic	76	49	12
Aspermic	2	14	18
Oligozoospermic	11	6	16
Severe oligozoospermic	8	4	4
Teratozoospermic	24	16	24
Asthenozoospermic	5	9	6
Total	126	98	80
Sperm motility			
Nonmotile	5	9	6
Slow progressive motile	40	35	38
Rapid progressive motile	71	36	14
Total	116	80	58
Sperm morphology			
Normal	76	36	15
Head defect	12	7	22
Tail defect	11	9	7
Multiple defect	17	28	14
Total	116	80	58
Sperm DNA integrity			
Normal (> 50%)	88	46	33
Damaged DNA	28	34	25
Total	116	80	57

Since the levels in both the smoking habit and motility categories can be ordered, we carried out ordinal logistic regression analysis. The frequency χ^2 -test shows significant association between the smoking habit and sperm motility ($\chi^2 = 23.188$, $p \approx 0.000$) (Table 3B). Further, the negative coefficient for smoking habit and odds ratio less than 1 indicates that higher smoking levels tend to be associated with lower motility levels.

The frequency χ^2 test shows significant association between smoking habit and sperm morphology ($\chi^2 = 25.90$ $p = 0.000$, Table 3B). Since morphology has two categories, normal and abnormal, binary logistic regression analysis was used to study the nature of association. Further, a positive coefficient for smoking habit and odds ratio greater than 1 indicates that higher smoking levels tend to be associated

Table 1. Demographic status of the population included in the present study.

Parameters	Non smokers (n = 126)	Light smokers (n = 98)	Heavy smokers (n = 80)	P value		
				NS/LS	NS/HS	LS/HS
Age (Mean \pm SD)	35 \pm 2.1	40 \pm 5.2	41 \pm 1	0.001	0.001	0.0921
BMI	25.92 \pm 4.77	25.68 \pm 5.92	21.31 \pm 4.22	0.737	0.001	0.001
No. of cigarette/day	0	1 - 20	> 21			
Sperm count $\times 10^6$ /ml	130 \pm 50	76 \pm 50	50 \pm 20	0.001	0.001	0.001

NS: Non smoker

LS: Light smoker

HS: Heavy smoker

Table 3A. Logistic regression analysis of the confounding effect of age and BMI on seminal phenotype and smoking.

Predictor	Coef	OR (95%CI)	P value
Logit 1: (5/0)			
Smoking	0.7091	2.03(1.04-3.98)	0.038
Age	-0.00471	1.00(0.89 - 1.12)	0.936
BMI	-0.14478	0.87(0.75 - 1.00)	0.050
Logit 2: (4/0)			
Smoking	0.7844	2.19(1.42 - 3.38)	0.000
Age	0.00563	1.01(0.93 - 1.08)	0.881
BMI	0.00520	1.01(0.92 - 1.10)	0.907
Logit 3: (3/0)			
Smoking	0.1020	1.11(0.53 - 2.33)	0.788
Age	0.06141	1.04(0.92 - 1.17)	0.528
BMI	0.07929	0.89(0.76 - 1.04)	0.142
Logit 4: (2/0)			
Smoking	1.0448	2.84(1.64 - 4.92)	0.000
Age	-0.00482	1.00(0.91 - 1.09)	0.918
BMI	0.00530	1.01(0.90 - 1.12)	0.924
Logit 5: (1/0)			
Smoking	1.4228	4.15(2.26 - 7.63)	0.000
Age	-0.01342	0.99(0.90 - 1.08)	0.781
BMI	-0.17059	0.84(0.74 - 0.95)	0.007

Phenotypes: 0(Reference Event)normozoospermic, 5 asthenozoospermic, 4 teratozoospermic, 3 severe oligozoospermic, 2 oligozoospermic, 1 azoozoospermic

with abnormal morphology. It was noted that among heavy smokers with abnormal morphology, the percentage having only a head defect is significantly greater than the percentage having only a tail defect ($p = 0.037$).

A significant association was also observed between smoking habit and DNA integrity ($\chi^2 = 9.704$, $p = 0.008$). As DNA integrity has only two categories, normal DNA and damaged DNA (Fig. 1), a binary logistic regression analysis was pursued. Further, a positive coefficient for smoking habit and an odds ratio greater than 1 indicates that a higher level of smoking tends to be associated with sperm DNA damaged.

A significant difference in the FSH, LH, and testosterone levels was observed between non-smokers and heavy smokers (Fig. 2). Testosterone levels were found to be significantly lower among heavy smoking-severe oligo/azoospermic patients as compared to nonsmoking-normozoospermic controls ($P < 0.0001$, Student's paired t test), while FSH and LH levels were significantly higher among heavy smoking-severe oligo/azoospermic patients (Fig. 2).

Among nonsmoking infertile men, four different CAG repeat sizes (alleles) ranging from 14 - 29 were observed (Fig. 3), and among heavy smoking-severe oligo/




Figure 1. Assessment of human sperm DNA integrity by acridine orange staining (40X). DNA integrity was assessed using acridine orange staining following sperm fixation. The percentage of green (normal DNA integrity) and orange red (abnormal DNA integrity) spermatozoa per 200 spermatozoa in each sample was calculated. An abnormal integrity of sperm DNA was considered as more than 30% denaturation (orange-red).

azoospermic study patients six different CAG repeat sizes were observed ranging from 16 - 33 (Table 4). The mean number of CAG repeats in the non-smoking group was $19.1 + 5.04$. In the heavy smoking group the repeat size was significantly higher $26 + 5.8$ ($p = 0.03$).

Despite worldwide anti-smoking campaigns, cigarette smoking is very common. The outcome of the present study should be analyzed in the context of the urban lifestyle of the city of Kolkata. However, the results reflect the changing semen quality of smokers attending an andrology laboratory with infertility related problems and should not be linked to the condition of the general fertile population. The subjects in the present study consisted of patients without extreme pathological conditions and had a sperm count $> 20 \times 10^6$ /ml. The study presented above shows that 58.5% of the smokers were among the group of infertile men. Moreover, the present study includes semen parameters with different seminal phenotypes, motility, morphology, DNA integrity, and level of endocrine hormones such as FSH, LH, and testosterone as well as androgen receptor genes with varied sized CAG repeats. Cigarette smoke is a

Table 3B. Logistic regression analysis of sperm motility, morphology, and DNA integrity with smoking habit.

Logit/Parameter	Predictor	Coeff.	p-value	OR (95%CI)
Sperm motility	Constant Smoking habit	-1.1493 -0.7045	0.000 0.000	0.49(0.36 - 0.68)
Sperm morphology	Constant Smoking habit	-1.4897 0.8467	-0.000 0.000	2.33 (1.66-3.27)
DNA integrity	Constant Smoking habit	-1.4939 0.4638	0.000 0.008	1.59 (1.15-2.21)

Reference level: normo




Figure 2. FSH, LH, and testosterone levels among non-smokers and smokers. Follicle stimulating hormone (FSH), leutanising hormone (LH), and testosterone concentrations were determined in the serum of smokers (severeoligo/azoospermic) and nonsmokers (normozoospermic). Values are mean and SD from two independent experiments. mIU, mili International Unit




Figure 3. CAG repeat length in the AR genes of subjects. The CAG repeat segment of the AR gene was amplified by PCR with primers and conditions described in Materials and Methods. The representative products analyzed by agarose gel electrophoresis are shown. L1 and L2 represent repeat numbers 33 and 16, respectively. M, 500 base pair DNA size marker; AR: androgen receptor

cell mutagen and carcinogen and certain components in cigarette smoke interact directly or indirectly with the male and female gametes affecting their function [Zenzes 2000]. Tobacco consumption involves about 4,000 compounds. The major components such as lead, cadmium, nicotine, and benzopyrene are known to affect semen parameters and sperm function [Zenzes 2000] as well as the endocrine status.

It is noted in this study that smoking is associated with decreased sperm concentration and motility (Table 2). The effects of tobacco smoking on reproductive function have been performed on a group of infertile patients. A large study conducted by Pasqualotto et al. [2006] showed a decline in semen volume with an increased number of cigarettes smoked, although no statistically significant differences were observed between the groups in terms of

sperm motility, concentration, or morphology [Zenzes 2000]. Another study reported a non-significant change towards higher sperm concentration with more smoking [Olderied et al. 1989], while others showed either a statistically significant dose response relation or a trend towards decreased sperm concentration with increased smoking [Zenzes 2000]. In heavy smokers, azoospermic / oligozoospermic / severe oligozoospermic / teratozoospermic / asthenozoospermic phenotypes were found in 85% of cases compared to 39% for non-smokers and 50% of smokers suggesting a trend towards deviation from normozoospermic phenotype with increased smoking. Earlier studies have concluded that toxins in cigarette smoke reach the male reproductive system and interact directly with components of the seminal fluid and the accessory endocrine system leading to increased viscosity, reduced seminal volume, and delayed liquefaction. This ultimately leads to the reduction of forward linear progression of spermatozoa, manifest as asthenozoospermia [Chia et al. 1994; Kulikauskas et al. 1985; Kunzle et al. 2003; Zhang et al. 2000]. However, in this population of infertile men, there is a trend towards increased oligozoospermic, teratozoospermic, and azoospermic phenotypes. The presence of the dominant asthenozoospermic phenotype has been observed in both smokers and non-smokers in the study conducted at Dehradun, India [Gaur et al. 2007].

Nicotine, present in the smoke, proved to be a potential oxidant which can affect spermatogenesis. Its damage to membrane structure of the spermatozoa is a major cause of the impaired motility [Tejada et al. 1984]. The mammalian sperm and neuron appear to share many of the 'neuronal' receptors. Exocytosis in neurons and sperm is essential to the functions of these cells and is strongly influenced by similar receptors. 'Neuronal' receptor types in sperm may also play a role in the control of sperm motility [Arabi 2004]. It is highly likely that nicotine disrupts the activity of the nicotinic acetylcholine receptor of spermatozoa and thus weakens the intracellular Ca^{+2} dynamics resulting in impaired motility.

The study showed poor sperm morphology among the smokers. Cigarettes contain heavy metals, which cause high oxidative stress. Due to the oxidative stress, membrane damage can occur which can lead to morphological defects. Heavy metals also cause DNA damage during spermatogenesis which may result in abnormal development of sperm topography. It was noted that DNA damage is increased with habitual smoking. There have been previous studies highlighting the induction of DNA damage as well as the inhibition of DNA repair mechanisms by heavy metals [Hengstler et al. 2003; Meizel 2004]. A range of molecular mechanisms was proposed, reflective of their diverse chemical properties. Heavy metals present in cigarette smoke, like lead and cadmium causes an increase in the production of reactive oxygen species (ROS) such as hydroxyl radical ($HO\cdot$), superoxide radical ($O_2^{\cdot-}$), or hydrogen peroxide (H_2O_2). Enhanced generation of ROS can overwhelm the cells' intrinsic antioxidant defenses, and result in 'oxidative stress' [Meizel 2004]. Cells under oxidative stress can cause

Table 4. Distribution of length of CAG repeats in the androgen receptor gene.

ID	Age	BMI (kg/m ²)	FSH (mIU/ml)	LH (mIU/ml)	Testosterone (ng/ml)	CAG Repeat in AR gene
N287	34	22.2	5.8	5.2	5.6	14
N40	40	23.1	7.2	6.8	5.2	22
N7	39	19.8	6.8	7.7	6.2	22
N24	38	25.5	7.7	6.2	4.2	29
N160	36	24.5	5.5	5.8	6.2	22
N304	38	26.7	7.4	6.6	5.5	22
N251	41	22.1	7.2	5.8	6.7	16
N216	35	19.9	6.5	6.6	5.6	16
N139	36	22.2	8.2	7.8	4.8	14
N128	40	20.5	5.6	5.2	5.4	14
Mean ± SD	37.7 ± 2.35	22.6 ± 2.3	6.79 ± 0.9	6.37 ± 0.9	5.54 ± 0.7	19.1 ± 5.04
A70	39	20.1	10	9.8	1.6	29
A82	42	17.6	11.5	10.6	1.9	22
A8	40	20.2	12.3	10.2	1.5	29
A30	36	19.2	12	11.2	3.2	16
A146	39	19.7	10.6	9.2	1.0	33
A113	42	20.4	11.2	10.6	2.2	22
A238	39	23.1	11.5	10.5	1.5	32
A298	43	28.5	9.8	7.8	1.2	29
A171	34	18.5	10.4	8.8	1.8	29
A195	41	20.2	10.5	7.8	2.0	19
Mean ± SD	39.5 ± 2.7	20.75 ± 3.07	10.98 ± 0.84	9.65 ± 1.2	1.79 ± 0.6	26 ± 5.8
			<i>P</i> < 0.0001	<i>P</i> < 0.0001	<i>P</i> < 0.0001	<i>P</i> = 0.033

N: non-smoking group

A: smoking group

oxidative DNA damage, strand breaks, and induce mutations [Hengstler et al. 2003]. A decrease in repair capacity may increase susceptibility to reactive oxygen species generated by these heavy metals [Hengstler et al. 2003]. It has also been shown that repair of 8-hydroxy-2'-deoxyguanosine decreased with increasing cadmium exposure and was inversely correlated with the level of DNA-single strand breaks [Meizel 2004]. In this study DNA fragmentation was estimated by acridine orange (AO) staining. In the AO test, the fluorescent dye intercalates between the bases in intact double-stranded DNA producing green fluorescence (530 nm). However, when there is a strand break, the dye forms a weaker bond with the single strand along the back-bone giving red fluorescence (650 nm) (Fig. 1). Since an AO-test can be done much more easily than tests like SCSA, it can serve as a routine screening method to estimate the gross DNA damage in a sperm population [Ercal et al. 2001].

The present study showed higher FSH, LH, and lower testosterone levels with increased smoking. This is in contrast to earlier studies which reported higher testosterone and LH levels in addition to higher LH/free testosterone ratios with increased smoking [Trummer et al. 2002], or a positive association between smoking and testosterone [Alexandersen et al. 1996]. Nicotine, one of the toxic components in cigarette smoke, had been reported to inhibit androgen biosynthesis and Leydig-cell growth [Bergmann et al. 1994; Funabashi et al. 2005] which might be responsible for the observed decline in testosterone among the heavy smokers. A high FSH level is usually diagnostic of primary testicular failure, a condition in which the seminiferous tubules in the testes do not produce sperm normally, because they are damaged.

Patients having mutations in AR gene exhibited elevated LH levels [Quigley 1998] which might be in response to the impaired feedback signal of testosterone in the pituitary. Leydig cell androgen secretion is mainly regulated via LH receptor mediated events and a receptor down regulation occurs in the presence of elevated LH levels. The mechanisms by which androgens regulate spermatogenesis are still elusive. In human testis, AR is expressed in Leydig as well as in peritubular and sertoli cells and provides sites for regulation of spermatogenesis and sperm outlet. The AR has a dual role in impaired spermatogenesis and reducing the levels of testosterone reflective of its CAG repeat length. The observed effect that CAG repeat length is positively correlated with smoking and inversely correlated with sperm concentration indicating that the variable polyglutamine tract size as deduced from the CAG repeat length in the AR gene, mediates androgen action with varying intensity. This modulation of AR androgenicity might be reflected by the serum testosterone level.

The above two studies [Davis-Dao et al. 2007; Mifsud et al. 2001] has shown a direct correlation of AR CAG length with free testosterone in fertile men. Although not observed in all populations [Dadze et al. 2000], depressed spermatogenesis has been associated with long CAG tracts in men from Singapore [Tut et al. 1997], Australia [Dowsing et al. 1999], Japan [Yoshida et al. 1999], and the USA [Mifsud et al. 2001; Patrizio et al. 2001]. A previous study by Thangaraj et al. [2002] shows no association of AR-CAG length with infertility when 280 azoospermic men were compared with 201 fertile men from this city. In the present study, we have included 20 infertile men out of which 10 were heavy smokers with low testosterone and 10

non-smokers with a normal level of testosterone and observed the AR-CAG length variation to assess the confounding effect of low testosterone and smoking. This study presents a significantly increased CAG repeat length among heavy smokers with low testosterone. Strikingly contradicting results exist in the literature regarding the association between testosterone and AR-CAG tract in subfertile patients [Mifsud et al. 2001]. However most of the studies have compared fertile and infertile men, and in this manner have tried to negate this bias and to observe the individual effect of smoking on CAG repeat length. Only infertile men were considered in the present study. Further studies on the effect of increased CAG repeat and smoking status with a larger sample size are required.

Materials and Methods

Sample population

A total of 304 men of Indian nationals around Kolkata city, who attended the out patient department of the Institute of Reproductive Medicine, Kolkata, India with a history of infertility for at least 1 year, and were able to provide an ejaculate were evaluated. Men meeting the following inclusion criteria were chosen: no vasectomy, no medical history of genital disease, and no abnormality in the reproductive organs. Subjects were excluded from the study if they had pathologies that aggravate erythrocytosis, cardiovascular, and renal disease. Full data was collected through a questionnaire including the number of cigarettes per day. For body mass index (BMI), height, and weight were obtained for each subject and BMI was calculated as kg/m^2 . Men who had smoked cigarettes for >6 months and were still smoking were classified as smokers. Smokers were categorized as light (1–20 cigarettes per day) and heavy (> 20 cigarettes per day) smokers. The study was approved by the institutional ethical committee as per guidelines.

Semen analyses

Semen samples were collected by masturbation in a sterile wide-mouthed plastic container after a sexual abstinence of 4–5 d. The sample was allowed to liquefy at 37° C for 20 min before analysis. Sperm concentration and motility were evaluated using a Makler's counting chamber (Sefi Medical Instruments, Haifa, Israel). Total sperm count of a given semen sample was calculated by multiplying the seminal sperm concentration with the volume. Sperm morphology was assessed using Papanicolaou staining method.

Semen analysis was carried out by standard WHO procedure as described earlier and was classified according to the nomenclature of semen variables [WHO 1999]. Normozoospermia was diagnosed when sperm concentration, motility, and morphology were within the reference values. The reference values for sperm count is $\geq 20 \times 10^6$ sperm/ml, for motility 50% sperm with forward progression, for morphology $\geq 30\%$ sperm with normal morphology, and for DNA integrity 50% sperm should have intact DNA. Oligozoospermia was determined when the sperm count was less than the reference value and severe-oligozoospermia

when sperm concentration was $< 5 \times 10^6$ /ml. Likewise, asthenozoospermia and teratozoospermia were diagnosed when motility and morphology were below the reference values respectively. Azoospermia was diagnosed when not a single spermatozoa was found in the ejaculate.

Sperm DNA integrity by acridine orange (AO) fluorescence

A smear was drawn on a clean slide using 20 μl of the semen sample. For assessment of sperm DNA integrity, the smears were air dried for 1 hr and then fixed overnight in freshly prepared Carnoy's solution (1:3 glacial acetic acid: methanol). The slides were washed in distilled water and allowed to dry before staining. Each sample was then stained in acridine orange (Sigma Chemical Co, St. Louis, USA) solution (10ml of 1% AO in distilled water added to a mixture of 40ml of 0.1 M citric acid and 2.5ml of 0.3 M $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$) for 5 min in dark. The preparations were gently washed in distilled water and examined on the same day using a Fluorescent microscope (Leica DM 3000, Leica Microsystem Ltd., Switzerland) excitation at 450 to 490 nm. The percentage of green (normal DNA integrity) and orange red (abnormal DNA integrity) spermatozoa per 200 spermatozoa in each sample was calculated by the same single person. An abnormal integrity of sperm DNA was considered as more than 30% denaturation (orange-red).

Hormone assays

Blood (~ 2 ml) was collected from the antecubital vein in vials without EDTA (clot vials) and serum isolated by centrifugation (1000 x rpm, 10 min). Follicle stimulating hormone (FSH), leutinising hormone (LH), and testosterone concentrations were determined in the serum using standard ELISA kits (Biorad Laboratories, USA) to evaluate the endocrine status of individuals involved in the study group consisting of heavy smoking-severeoligo/azoospermic patients while nonsmoking-normozoospermic group were kept as controls.

DNA isolation

Genomic DNA was extracted from peripheral blood of heavy-smoker (with low testosterone level) and non-smoker subjects using a standard procedure. Briefly, erythrocytes were lysed using erythrocyte lysis buffer (10 mM Tris pH 8.0, 320 mM sucrose, 5 mM MgCl_2 , 1% Triton X - 100) for 5 min. Leukocytes were pelleted by centrifugation at 500 x g for 5 min, and the pellet dissolved in leukocyte lysis buffer (400mM Tris, 60mM EDTA, 150 mM NaCl, 1% SDS). 5M sodium perchlorate (Sigma Aldrich, USA) was added and mixed thoroughly for 2–3min. DNA was extracted with chloroform and precipitated with isopropanol and finally washed with 70% ethanol and dissolved in TE buffer (10 mM Tris pH 8, 1 mM EDTA).

PCR assay for CAG repeat count

CAG repeat segment in exon 1 of androgen receptor (AR) gene was amplified with the primers: forward 5'AGAGCGTGCGG-GAAGCGTTCCAGAACCCG3' and reverse 3'AACGTG-GATGGGGCAGCTGAGTCAT5'. PCR amplification of

each sample was performed using 5 ng of template DNA 10 pM of each primer, 2.5 mM deoxy Nucleotide triphosphate, 1X PCR buffer containing 1.5mM MgCl₂ and 0.5 u of Taq polymerase enzyme (Bangalore Genei, India). Amplification was carried out in ABI 2700 thermal cycler (Applied Biosystems, USA) using the following conditions: 30 cycles of denaturation at 94°C for 30 sec, annealing at 55°C for 45 sec, and extension at 72°C for 60 sec. The final extension was at 72°C. In the first cycle the sample was denatured for 5min.

Automated DNA sequencing

To confirm CAG repeat numbers duplicate samples from each repeat size were sequenced in ABI PRISM 3700 (Perkin Elmer, USA) automated DNA sequencers using the manufacturers protocol.

Statistical analysis

Statistical analysis of the data has been performed using MINITAB™ Statistical Software (version: 13.31, Minitab Inc. PA, USA). Frequency χ^2 -test has been used to examine the association between smoking habit and seminal phenotype, sperm motility, morphology, and DNA integrity. Logistic regression analysis was carried out to study the nature of dependence, wherever applicable. A *p*-value less than 0.05 were considered as statistically significant.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Author Contributions: Amrita Mitra: Literature search, experiments, data-collection and manuscript preparation
Baidyanath Chakraborty: Study designing and clinical evaluation

Dyutiman Mukhopadhyay: Data-analysis

Manisha Pal: Statistical analysis

Sanjit Mukherjee: manuscript preparation and experimentation

Samir Banerjee: Study designing and data-analysis

Keya Chaudhuri: Literature search, study design, data analysis, data interpretation, manuscript preparation

References

- Alexandersen, P., Haarbo, J. and Christiansen, C. (1996) The relationship of natural androgens to coronary heart disease in males: a review. *Atherosclerosis* **125**:1–13.
- Arabi, M. (2004) Nicotinic infertility: assessing DNA and plasma membrane integrity of human spermatozoa. *Andrologia* **36**:305–310.
- Bergmann, M., Behre, H.M. and Nieschlag, E. (1994) Serum FSH and testicular morphology in male infertility. *Clin Endocrinol (Oxf)* **40**:133–136.
- Chang, C.S., Kokontis, J. and Liao, S.T. (1988) Molecular cloning of human and rat complementary DNA encoding androgen receptors. *Science* **240**:324–326.
- Chia, S.E., Ong, C.N. and Tsakok, F.M. (1994) Effects of cigarette smoking on human semen quality. *Arch Androl* **33**:163–168.
- Dadze, S., Wieland, C., Jakubiczka, S., Funke, K., Schroder, E., Royer-Pokora, B., et al. (2000) The size of the CAG repeat in exon 1 of

- the androgen receptor gene shows no significant relationship to impaired spermatogenesis in an infertile Caucasoïd sample of German origin. *Mol Hum Reprod* **6**:207–214.
- Davis-Dao, C.A., Tuazon, E.D., Sokol, R.Z. and Cortessis, V.K. (2007) Male infertility and variation in CAG repeat length in the androgen receptor gene: a meta-analysis. *J Clin Endocrinol Metab* **92**:4319–4326.
- Dowsing, A.T., Yong, E.L., Clark, M., McLachlan, R.I., de Kretser, D. M. and Trounson, A.O. (1999) Linkage between male infertility and trinucleotide repeat expansion in the androgen-receptor gene. *Lancet* **354**:640–643.
- Ercal, N., Gurer-Orhan, H. and Aykin-Burns, N. (2001) Toxic metals and oxidative stress part I: mechanisms involved in metal-induced oxidative damage. *Curr Top Med Chem* **1**:529–539.
- Funabashi, T., Sano, A., Mitsushima, D. and Kimura, F. (2005) Nicotine inhibits pulsatile luteinizing hormone secretion in human males but not in human females, and tolerance to this nicotine effect is lost within one week of quitting smoking. *J Clin Endocrinol Metab* **90**:3908–3913.
- Gaur, D.S., Talekar, M. and Pathak, V.P. (2007) Effect of cigarette smoking on semen quality of infertile men. *Singapore Med J* **48**:119–123.
- Hengstler, J.G., Bolm-Audorff, U., Faldum, A., Janssen, K., Reifenrath, M., Gotte, W., et al. et al (2003) Occupational exposure to heavy metals: DNA damage induction and DNA repair inhibition prove co-exposures to cadmium, cobalt and lead as more dangerous than hitherto expected. *Carcinogenesis* **24**:63–73.
- Kulikauskas, V., Blaustein, D. and Ablin, R.J. (1985) Cigarette smoking and its possible effects on sperm. *Fertil Steril* **44**:526–528.
- Kunzle, R., Mueller, M.D., Hanggi, W., Birkhauser, M.H., Drescher, H. and Bersinger, N.A. (2003) Semen quality of male smokers and nonsmokers in infertile couples. *Fertil Steril* **79**:287–291.
- Lo, K.C., Brugh, V.M., 3rd, Parker, M., and Lamb, D.J. (2005) Isolation and enrichment of murine spermatogonial stem cells using rhodamine 123 mitochondrial dye. *Biol Reprod* **72**:767–771.
- Meizel, S. (2004) The sperm, a neuron with a tail: 'neuronal' receptors in mammalian sperm. *Biol Rev Camb Philos Soc* **79**:713–732.
- Mifsud, A., Choon, A.T., Fang, D. and Yong, E.L. (2001) Prostate-specific antigen, testosterone, sex-hormone binding globulin and androgen receptor CAG repeat polymorphisms in subfertile and normal men. *Mol Hum Reprod* **7**:1007–1013.
- Mifsud, A., Sim, C.K., Boettger-Tong, H., Moreira, S., Lamb, D.J., Lipschultz, L.I., et al. (2001) Trinucleotide (CAG) repeat polymorphisms in the androgen receptor gene: molecular markers of risk for male infertility. *Fertil Steril* **75**:275–281.
- Mukhopadhyay, D., Nandi, P., Varghese, A.C., Gutgutia, R., Banerjee, S. and Bhattacharyya, A.K. (2010) The in vitro effect of benzo[a]pyrene on human sperm hyperactivation and acrosome reaction. *Fertil Steril* **94**:595–598.
- Oldereid, N.B., Rui, H., Clausen, O.P. and Purvis, K. (1989) Cigarette smoking and human sperm quality assessed by laser-Doppler spectroscopy and DNA flow cytometry. *J Reprod Fertil* **86**:731–736.
- Padron, R.S., Mas, J., Zamora, R., Riverol, F., Licea, M., Mallea, L., et al. (1989) Lipids and testicular function. *Int Urol Nephrol* **21**:515–519.
- Pasqualotto, F.F., Sobreiro, B.P., Hallak, J., Pasqualotto, E.B. and Lucon, A.M. (2006) Cigarette smoking is related to a decrease in semen volume in a population of fertile men. *BJU Int* **97**:324–326.
- Patrizio, P., Leonard, D.G., Chen, K.L., Hernandez-Ayup, S. and Trounson, A.O. (2001) Larger trinucleotide repeat size in the androgen receptor gene of infertile men with extremely severe oligozoospermia. *J Androl* **22**:444–448.
- Quigley, C.A. (1998) The androgen receptor: physiology and pathophysiology. In: Nieschlag E, Behre HM, eds. *Testosterone: action, deficiency, substitution*. New York: Springer; 33–106
- Tejada, R.I., Mitchell, J.C., Norman, A., Marik, J.J. and Friedman, S. (1984) A test for the practical evaluation of male fertility by acridine orange (AO) fluorescence. *Fertil Steril* **42**:87–91.
- Thangaraj, K., Joshi, M.B., Reddy, A.G., Gupta, N.J., Chakravarty, B. and Singh, L. (2002) CAG repeat expansion in the androgen

- receptor gene is not associated with male infertility in Indian populations. *J Androl* **23**:815–818.
- Trisini, A.T., Singh, N.P., Duty, S.M. and Hauser, R. (2004) Relationship between human semen parameters and deoxyribonucleic acid damage assessed by the neutral comet assay. *Fertil Steril* **82**:1623–1632.
- Trummer, H., Habermann, H., Haas, J. and Pummer, K. (2002) The impact of cigarette smoking on human semen parameters and hormones. *Hum Reprod* **17**:1554–1559.
- Tut, T.G., Ghadessy, F.J., Trifiro, M.A., Pinsky, L. and Yong, E.L. (1997) Long polyglutamine tracts in the androgen receptor are associated with reduced trans-activation, impaired sperm production, and male infertility. *J Clin Endocrinol Metab* **82**:3777–3782.
- Varghese, A.C., Bragais, F.M., Mukhopadhyay, D., Kundu, S., Pal, M., Bhattacharyya, A.K., et al. (2009) Human sperm DNA integrity in normal and abnormal semen samples and its correlation with sperm characteristics. *Andrologia* **41**:207–215.
- WHO (1999) WHO laboratory manual for the examination of Human Semen and Sperm-Cervical Mucus Interaction, 4th edn, Cambridge University Press, Cambridge, UK.
- Yong, E.L., Ghadessy, F., Wang, Q., Mifsud, A. and Ng, S.C. (1998) Androgen receptor transactivation domain and control of spermatogenesis. *Rev Reprod* **3**:141–144.
- Yoshida, K.I., Yano, M., Chiba, K., Honda, M. and Kitahara, S. (1999) CAG repeat length in the androgen receptor gene is enhanced in patients with idiopathic azoospermia. *Urology* **54**:1078–1081.
- Zenzen, M.T. (2000) Smoking and reproduction: gene damage to human gametes and embryos. *Hum Reprod Update* **6**:122–131.
- Zhang, J.P., Meng, Q.Y., Wang, Q., Zhang, L.J., Mao, Y.L. and Sun, Z. X. (2000) Effect of smoking on semen quality of infertile men in Shandong, China. *Asian J Androl Jun* **2**(2):143–6